

# K3 Surfaces and Lattice Theory

Ichiro Shimada

Hiroshima University

2014 Dec: Hanoi

## Example

Consider two surfaces  $S_+$  and  $S_-$  in  $\mathbb{C}^3$  defined by

$$w^2(G(x, y) \pm \sqrt{5} \cdot H(x, y)) = 1, \quad \text{where}$$

$$G(x, y) := -9x^4 - 14x^3y + 58x^3 - 48x^2y^2 - 64x^2y \\ + 10x^2 + 108xy^3 - 20xy^2 - 44y^5 + 10y^4,$$

$$H(x, y) := 5x^4 + 10x^3y - 30x^3 + 30x^2y^2 + \\ + 20x^2y - 40xy^3 + 20y^5.$$

Since  $S_+$  and  $S_-$  are conjugate by  $\text{Gal}(\mathbb{Q}(\sqrt{5})/\mathbb{Q})$ , they can *not* be distinguished algebraically.

But  $S_+$  and  $S_-$  are not homeomorphic (in the classical topology).

Many examples of non-homeomorphic conjugate complex varieties are known since Serre (1964).

# Introduction

## Definition

A smooth projective surface  $X$  is called a *K3 surface* if

- $\exists$  a nowhere vanishing holomorphic 2-form  $\omega_X$  on  $X$ , and
- $\pi_1(X) = \{1\}$ .

We consider the following geometric problems on  $K3$  surfaces:

- enumerate elliptic fibrations on a given  $K3$  surface,
- enumerate elliptic  $K3$  surfaces up to some equivalence relation,
- enumerate projective models of a given  $K3$  surface,
- enumerate projective models of  $K3$  surfaces,
- determine the automorphism group of a given  $K3$  surface,
- .....

## The aim of this talk

Thanks to the theory of period mapping for  $K3$  surfaces and the Torelli-type theorem due to Piatetski-Shapiro and Shafarevich, some of these problems are reduced to computational problems in lattice theory, and the latter can often be solved by means of *computer*.

In this talk, we explain how to use lattice theory and computer in the study of  $K3$  surfaces.

We then demonstrate this method on the problems of constructing Zariski pairs of plane curves of degree 6.

A *lattice* is a free  $\mathbb{Z}$ -module  $L$  of finite rank with a non-degenerate symmetric bilinear form

$$\langle \quad \rangle: L \times L \rightarrow \mathbb{Z}.$$

Let  $L$  be a lattice of rank  $n$ . We choose a basis  $e_1, \dots, e_n$  of  $L$ . The lattice  $L$  is given by the Gram matrix

$$G := (\langle e_i, e_j \rangle)_{i,j=1,\dots,n}.$$

- $O(L)$  is the group of all isometries of  $L$ .
- $L$  is *unimodular* if  $\det G = \pm 1$ .
- The *signature*  $\text{sgn}(L)$  is the signature of the real quadratic space  $L \otimes \mathbb{R}$ .
- A lattice  $L$  is said to be *hyperbolic* if  $\text{sgn}(L) = (1, n - 1)$ , and is *positive-definite* if  $\text{sgn}(L) = (n, 0)$ .
- A lattice  $L$  is *even* if  $v^2 \in 2\mathbb{Z}$  for all  $v \in L$ .
- A sublattice  $L'$  of  $L$  is *primitive* if  $L/L'$  is torsion free.

## Lattices associated to a $K3$ surface

$K3$  surfaces are diffeomorphic to each other.

Suppose that  $X$  is a  $K3$  surface.

Then  $H^2(X, \mathbb{Z})$  with the cup product is an even unimodular lattice of signature  $(3, 19)$ , and hence is isomorphic to the  $K3$  lattice

$$U^{\oplus 3} \oplus E_8^{-\oplus 2},$$

where  $U$  is the hyperbolic plane with a Gram matrix

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

and  $E_8^-$  is the negative definite root lattice of type  $E_8$ .

$$\begin{bmatrix} -2 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & -2 \end{bmatrix}$$

The Gram matrix of  $E_8^-$

## The Néron-Severi lattice

$$S_X := H^2(X, \mathbb{Z}) \cap H^{1,1}(X)$$

is the sublattice of  $H^2(X, \mathbb{Z})$  generated by classes of curves on  $X$ , which is primitive. It is an even hyperbolic lattice of rank  $\leq 20$ . Moreover the sublattice  $S_X$  of  $H^2(X, \mathbb{Z})$  is primitive.

Our goal is to extract geometric information of  $X$  from the Gram matrix of  $S_X$ .

### Problem

*Suppose that an even hyperbolic lattice  $S$  of rank  $\leq 20$  is given. Is there a K3 surface  $X$  such that  $S \cong S_X$ ?*



By the *surjectivity of the period map*, we have the following:

### Theorem

Let  $S$  be a primitive hyperbolic sublattice of  $U^{\oplus 3} \oplus E_8^{-\oplus 2}$ . Then there exists a K3 surface  $X$  such that  $S \cong S_X$ .

### Problem

Suppose that an even lattice  $L$  and an even unimodular lattice  $M$  are given. Can  $L$  be embedded into  $M$  primitively?

A lattice  $L$  is canonically embedded into its *dual lattice*

$$L^\vee := \text{Hom}(L, \mathbb{Z})$$

as a submodule of finite index. The finite abelian group

$$D_L := L^\vee / L$$

is called the *discriminant group* of  $L$ .

The symm. bil. form on  $L$  extends to a  $\mathbb{Q}$ -valued symm. bil. form on  $L^\vee$ , and it defines a finite quadratic form

$$q_L: D_L \rightarrow \mathbb{Q}/2\mathbb{Z}, \quad \bar{x} \mapsto x^2 \bmod 2\mathbb{Z}.$$

**The calculation of  $(D_L, q_L)$ .** Let  $G$  be a Gram matrix of  $L$ . We have  $U, V \in GL_n(\mathbb{Z})$  such that

$$VGU^{-1} = \begin{pmatrix} d_1 & & & \\ & \ddots & & \\ & & \ddots & \\ & & & d_n \end{pmatrix},$$

with  $1 = d_1 = \dots = d_k < d_{k+1} \leq \dots \leq d_n$ . Then

$$D_L \cong \bigoplus_{i>k} \mathbb{Z}/(d_i).$$

The  $i$ th row vector of  $U$ , regarded as an element of  $L^\vee$  with respect to the dual basis  $e_1^\vee, \dots, e_n^\vee$ , generate the factor  $\mathbb{Z}/(d_i)$  of  $D_L$ .

## Theorem (Hasse principle)

*Suppose that  $s_+, s_- \in \mathbb{Z}_{\geq 0}$  and a finite quadratic form  $(D, q)$  are given. We can determine by an effective method whether there exists an even lattice  $L$  such that  $\text{sgn}(L) = (s_+, s_-)$  and  $(D_L, q_L) \cong (D, q)$ .*

## Theorem

*Let  $M$  be an even unimodular lattice. We can see whether  $\exists$  a primitive embedding  $L \hookrightarrow M$  by seeing whether  $\exists$  the “orthogonal complement” of  $L$  in  $M$ , which is characterized by the signature and the discriminant form.*

## Corollary

*We can determine whether a given even hyperbolic lattice of rank  $\leq 20$  is a Néron–Severi lattice of a K3 surface  $X$  or not.*

## Polarized K3 surfaces

We consider the projective models of  $X$ . For  $h \in S_X \cong \text{Pic}(X)$ , let  $\mathcal{L}_h \rightarrow X$  be a line bundle whose class is  $h$ .

### Definition

A vector  $h \in S_X$  of  $h^2 = d > 0$  is a *polarization of degree  $d$*  if  $|\mathcal{L}_h| \neq \emptyset$  and has no fixed-components.

Let  $h$  be a polarization of degree  $d$ . Then  $|\mathcal{L}_h|$  defines  $\Phi_h : X \rightarrow \mathbb{P}^{1+d/2}$ . We denote by

$$X \longrightarrow X_h \longrightarrow \mathbb{P}^{1+d/2}$$

the Stein factorization of  $\Phi_h$ . The normal surface  $X_h$  is the *projective model* of  $(X, h)$ , and has only rational double points as its singularities.

## Example

A plane curve  $B \subset \mathbb{P}^2$  is a *simple sextic* if  $B$  is of degree 6 and has only simple singularities (*ADE*-singularities). Let  $B$  be a simple sextic, and  $Y_B \rightarrow \mathbb{P}^2$  the double covering branched along  $B$ . The minimal resolution  $X_B$  of  $Y_B$  is a  $K3$  surface.

We denote by

$$\Phi_B : X_B \rightarrow Y_B \rightarrow \mathbb{P}^2$$

the composite of the min. resol. and the double covering, and by  $h_B \in S_{X_B}$  the class of the pull-back of a line. Then  $h_B$  is a polarization of degree 2, and  $Y_B$  is its projective model.

## Problem

*Suppose that  $h \in S_X$  with  $h^2 > 0$  is given. Is  $h$  a polarization? If so, what is the ADE-type of  $\text{Sing } X_h$ ?*

We consider the second problem first. Suppose that  $h$  is a polarization.

## Proposition

*The ADE-type of  $\text{Sing } X_h$  is equal to the ADE-type of the root system  $\{r \in S_X \mid \langle h, r \rangle = 0, \langle r, r \rangle = -2\}$ .*

The sublattice  $\{x \in S_X \mid \langle h, x \rangle = 0\}$  is negative-definite.

## Problem

*Given a positive-definite lattice  $L$ . Calculate the set  $\{r \in L \mid \langle r, r \rangle = 2\}$ .*

For a triple  $QT := [Q, \lambda, c]$ , where

- $Q$  is a pos-def  $n \times n$  symmetric matrix with entries in  $\mathbb{Q}$ ,
- $\lambda$  is a column vector of length  $n$  with entries in  $\mathbb{Q}$ ,
- $c \in \mathbb{Q}$ ,

we define  $F_{QT} : \mathbb{R}^n \rightarrow \mathbb{R}$  by


$$F_{QT}(v) := v Q^t v + 2 v \lambda + c.$$

We have an algorithm to calculate the finite set

$$E(QT) := \{ v \in \mathbb{Z}^n \mid F_{QT}(v) \leq 0 \}.$$

### Corollary

*When a polarization  $h$  is given, we can determine the ADE-type of  $\text{Sing } X_h$ .*





Let  $L$  be an even hyperbolic lattice. Let  $\mathcal{P}_L$  be one of the two connected components of  $\{x \in L \otimes \mathbb{R} \mid x^2 > 0\}$ .

For  $v \in L \otimes \mathbb{R}$  with  $v^2 < 0$ , we put

$$(v)^\perp := \{x \in \mathcal{P}_L \mid \langle x, v \rangle = 0\}.$$

We put

$$\mathcal{R}_L := \{r \in L \mid r^2 = -2\}.$$

Each  $r \in \mathcal{R}_L$  defines a reflection  $s_r \in O(L)$  into  $(r)^\perp$ :

$$s_r : x \mapsto x + \langle x, r \rangle r.$$

The closure in  $\mathcal{P}_L$  of each connected component of

$$\mathcal{P}_L \setminus \bigcup_{r \in \mathcal{R}_L} (r)^\perp$$

is a standard fundamental domain of the action on  $\mathcal{P}_L$  of

$$W(L) := \langle s_r \mid r \in \mathcal{R}_L \rangle.$$

Let  $\mathcal{P}(X) \subset S_X \otimes \mathbb{R}$  be the positive cone that contains an ample class (e.g., the class of a hyperplane section).

### Proposition

*By Riemann-Roch, we see that the cone*

$$N(X) := \{x \in \mathcal{P}(X) \mid \langle x, [C] \rangle \geq 0 \text{ for any curve } C \text{ on } X \}.$$

*is a std. fund. domain of the action of  $W(S_X)$  on  $\mathcal{P}(X)$ .*

It is obvious that, if  $h$  is a polarization, then  $h \in N(X)$ . For the converse, we need an additional condition. For example,

### Proposition

*A vector  $h \in S_X$  with  $h^2 = 2$  is a polarization of degree 2 if and only if  $h \in N(X)$  and  $\{e \in S_X \mid e^2 = 0, \langle e, h \rangle = 1\} = \emptyset$ .*

## Problem

Suppose that  $h \in S_X$  with  $h^2 > 0$  is given.  
Does  $h$  belong to  $N(X)$ ?


When we have an ample vector  $h_0 \in N(X)$ , this problem is reduced to the following:

## Problem

Suppose that we are given vectors  $h_0, h \in \mathcal{P}_L$ . Calculate the set

$$\{ r \in L \mid \langle r, h_0 \rangle > 0, \langle r, h \rangle < 0, \langle r, r \rangle = -2 \}.$$

There is an algorithm for this task.



## Zariski pairs

For a simple sextic  $B \subset \mathbb{P}^2$ ,

- $R_B$  : the  $ADE$ -type of  $\text{Sing } B$ ,
- $\text{degs } B$  : the list of degrees of irreducible components of  $B$ .

We say that  $B$  and  $B'$  are of the same config type and write  $B \sim_{\text{cfg}} B'$  if

- $R_B = R_{B'}$ ,  $\text{degs } B = \text{degs } B'$ ,
- their intersection patterns of irreducible comps are same.

### Example

Zariski showed the existence of a pair  $[B, B']$  such that

- $R_B = R_{B'} = 6A_2$ ,  $\text{degs } B = \text{degs } B' = [6]$ , and
- $\pi_1(\mathbb{P}^2 \setminus B) \cong \mathbb{Z}/(2) * \mathbb{Z}/(3)$ ,  $\pi_1(\mathbb{P}^2 \setminus B') \cong \mathbb{Z}/(2) \times \mathbb{Z}/(3)$ .

For a simple sextic  $B$  with

$$\Phi_B : X_B \rightarrow Y_B \rightarrow \mathbb{P}^2,$$

let  $\mathcal{E}_B$  be the set of exceptional curves of  $X_B \rightarrow Y_B$ , and let

$$\Sigma_B := \langle [E] \mid E \in \mathcal{E}_B \rangle \oplus \langle h_B \rangle \subset S_{X_B} \subset H^2(X_B, \mathbb{Z}),$$

where  $h_B$  is the class of the pull-back of a line. We denote the primitive closure of  $\Sigma_B$  by

$$\overline{\Sigma}_B \subset S_{X_B} \subset H^2(X_B, \mathbb{Z}).$$

After the partial results by Urabe, Yang (1996) made the complete list configuration type of simple sextics by classifying all such  $\overline{\Sigma}_B$ , and found 11159 types.

We write  $B \sim_{\text{emb}} B'$  if there exists a homeomorphism

$$\psi : (\mathbb{P}^2, B) \xrightarrow{\sim} (\mathbb{P}^2, B').$$

We have  $B \sim_{\text{emb}} B' \implies B \sim_{\text{cfg}} B'$ .

$\#$  of config types = 11159 <  $\#$  of emb-top types = ?

### Definition

A *Zariski pair* is a pair  $[B, B']$  of simple sextics such that  $B \sim_{\text{cfg}} B'$  but  $B \not\sim_{\text{emb}} B'$ .

We consider the finite abelian group

$$G(B) := \overline{\Sigma}_B / \Sigma_B.$$

We put

$$\Theta_B := (\Sigma_B \subset H^2(X_B, \mathbb{Z}))^\perp.$$

### Theorem

*If  $B \sim_{\text{emb}} B'$ , then  $\Theta_B \cong \Theta_{B'}$ .*

In fact,  $\Theta_B$  is a topological invariant of the open surface

$$U_B := \Phi_B^{-1}(\mathbb{P}^2 \setminus B) \subset X_B,$$

because we have  $\Theta_B \cong H^2(U_B, \mathbb{Z}) / \text{Ker}$ , where

$$\text{Ker} := \{ v \in H^2(U_B) \mid \langle v, x \rangle = 0 \text{ for all } x \in H^2(U_B) \}.$$



Since  $\Theta_B^\perp = \overline{\Sigma}_B$ , the discriminant groups of  $\overline{\Sigma}_B$  and  $\Theta_B$  are isomorphic,

### Corollary

*If  $B \sim_{\text{cfg}} B'$  but  $|G(B)| \neq |G(B')|$ , then  $B \not\sim_{\text{emb}} B'$ .*

This corollary produces many examples of Zariski pairs.

### Example

In Zariski's example  $[B, B']$  with  $R_B = R_{B'} = 6A_2$ ,  $\text{degs } B = \text{degs } B' = [6]$  and

$$\pi_1(\mathbb{P}^2 \setminus B) \cong \mathbb{Z}/(2) * \mathbb{Z}/(3), \quad \pi_1(\mathbb{P}^2 \setminus B') \cong \mathbb{Z}/(2) \times \mathbb{Z}/(3),$$

we have  $G(B) \cong \mathbb{Z}/3\mathbb{Z}$  and  $G(B') = 0$ .

## Singular K3 surfaces

### Definition

A K3 surface  $X$  is called *singular* if  $\text{rank}(S_X) = 20$ .

### Theorem (Shioda and Inose)

The map

$$X \mapsto T(X) := (S_X \subset H^2(X, \mathbb{Z}))^\perp$$

is a bijection from the set of isom. classes of singular K3 surfaces to the set of isom. classes of oriented pos.-definite even lattices of rank 2.

In fact, Shioda and Inose gave a recipe to construct the singular K3 surface  $X$  from the lattice  $T(X)$ .

In particular, every singular K3 surface  $X$  is defined over  $\overline{\mathbb{Q}}$ , and a Gram matrix of  $S_X$  is always available.

## Theorem (S. and Schütt)

*Let  $X$  and  $X'$  be singular K3 surfaces defined over  $\overline{\mathbb{Q}}$  such that  $q_{T(X)} \cong q_{T(X')}$ . Then there exists  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  such that  $X' \cong X^\sigma$ .*

If  $B$  is a simple sextic with total Milnor number 19, then  $X_B$  is a singular K3 surface with  $\Theta_B \cong T(X_B)$ .

## Corollary

*Let  $B$  be a simple sextic with total Milnor number 19 defined over  $\overline{\mathbb{Q}}$ . If the genus containing  $T(X_B)$  contains more than one isom. class of lattices, then  $\exists \sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  such that  $B \not\sim_{\text{emb}} B^\sigma$ .*

Thus we obtain example of *arithmetic Zariski pairs*.

## The first example revisited

Consider the config type of sextics  $B = L + Q$ , where

- $\deg L = 1$ ,  $\deg Q = 5$ ,
- $L$  and  $Q$  are tangent at one point with multiplicity 5 ( $A_9$ -singularity), and
- $Q$  has one  $A_{10}$ -singular point.

Such sextics are projectively isomorphic to

$$z \cdot (G(x, y, z) \pm \sqrt{5} \cdot H(x, y, z)) = 0,$$

where  $G(x, y, z)$  and  $H(x, y, z)$  are homogenizations of the polynomials in the 1st slide with  $L = \{z = 0\}$ .

The genus containing  $T(X_B)$  consists of

$$\begin{bmatrix} 2 & 1 \\ 1 & 28 \end{bmatrix} \text{ (for } +\sqrt{5}\text{)}, \quad \begin{bmatrix} 8 & 3 \\ 3 & 8 \end{bmatrix} \text{ (for } -\sqrt{5}\text{)}.$$

## Example

Consider two surfaces  $S_+$  and  $S_-$  in  $\mathbb{C}^3$  defined by

$$w^2(G(x, y) \pm \sqrt{5} \cdot H(x, y)) = 1, \quad \text{where}$$

$$G(x, y) := -9x^4 - 14x^3y + 58x^3 - 48x^2y^2 - 64x^2y \\ + 10x^2 + 108xy^3 - 20xy^2 - 44y^5 + 10y^4,$$

$$H(x, y) := 5x^4 + 10x^3y - 30x^3 + 30x^2y^2 + \\ + 20x^2y - 40xy^3 + 20y^5.$$

Since  $S_+$  and  $S_-$  are conjugate by  $\text{Gal}(\mathbb{Q}(\sqrt{5})/\mathbb{Q})$ , they can *not* be distinguished algebraically.

But  $S_+$  and  $S_-$  are not homeomorphic (in the classical topology).

Many examples of non-homeomorphic conjugate complex varieties are known since Serre (1964).