

# RATIONAL DOUBLE POINTS ON ENRIQUES SURFACES

ICHIRO SHIMADA

ABSTRACT. We classify, up to some lattice-theoretic equivalence, all possible configurations of rational double points that can appear on a surface whose minimal resolution is a complex Enriques surface.

## 1. INTRODUCTION

We work over the complex number field  $\mathbb{C}$ .

The automorphism group of an Enriques surface changes in a complicated way under specializations of the surface ([3, Section 3.1], [24, Remark 7.17]), and smooth rational curves on the surface control the change of the automorphism group. Hence the study of configurations of smooth rational curves are important for the explicit description of variations of automorphism groups on the family of Enriques surfaces (see Nikulin [16]). On the other hand, in the investigation of  $\mathbb{Q}$ -homology projective planes, Hwang-Keum-Ohashi [11] and Schütt [18] classified all possible maximal root systems of smooth rational curves on Enriques surfaces.

In this paper, we classify all root systems of smooth rational curves on Enriques surfaces up to certain equivalence relation. The list we obtain (see Theorem 1.7 and Table 1.1) includes, of course, the 31 root systems of rank 9 classified in [11] and [18]. Our method is purely lattice-theoretic and algorithmic. The main tool is the generalized Borcherds algorithm ([5], [6], [22]) to calculate the orthogonal group of a hyperbolic lattice. An advantage of our method is that we can obtain the whole result by a single set of algorithms, and that we are exempted from the case-by-case investigation of possible root systems.

A lattice  $L$  of rank  $n$  is said to be *hyperbolic* if the real quadratic space  $L \otimes \mathbb{R}$  is of signature  $(1, n - 1)$ . Let  $L$  be a hyperbolic lattice. A *positive cone* of  $L$  is one of the two connected components of  $\{x \in L \otimes \mathbb{R} \mid \langle x, x \rangle > 0\}$ . Let  $O^+(L)$  denote the group of isometries of  $L$  that preserve a positive cone.


A *root* of a lattice  $L$  is a vector of square-norm  $-2$ . We say that  $L$  is a *root lattice* if  $L$  is generated by roots. An *ADE-configuration of roots* of  $L$  is a finite set  $\Phi$  of roots of  $L$  such that each connected component of the Dynkin diagram of  $\Phi$  is of type  $A_l$  ( $l \geq 1$ ),  $D_m$  ( $m \geq 4$ ), or  $E_n$  ( $n = 6, 7, 8$ ). The *ADE-type*  $\tau(\Phi)$  of an ADE-configuration  $\Phi$  of roots is the ADE-type of the Dynkin diagram of  $\Phi$ . It is well-known that a negative-definite root lattice  $R$  has an ADE-configuration  $\Phi_R$  of roots that form a basis of  $R$ . We define the ADE-type  $\tau(R)$  of  $R$  to be  $\tau(\Phi_R)$ . Conversely, for an ADE-type  $t$ , we have a negative-definite root lattice  $R(t)$ , unique up to isomorphism, such that  $\tau(R(t)) = t$ .

---

2010 *Mathematics Subject Classification.* 14J28, 14Q10.

*Key words and phrases.* Enriques surface, rational double points, hyperbolic lattice.

This work was supported by JSPS KAKENHI Grant Number 16H03926 and 16K13749.

FIGURE 1.1. Dynkin diagram of the basis  $E_{10}$  of  $L_{10}$ 

Let  $L_{10}$  be an even unimodular hyperbolic lattice of rank 10, which is unique up to isomorphism, and which has a basis  $E_{10} := \{e_1, \dots, e_{10}\}$  consisting of roots whose Dynkin diagram is given in Figure 1.1.

For a smooth projective surface  $Z$ , we denote by  $S_Z$  the lattice of numerical equivalence classes of divisors of  $Z$ , and by  $\mathcal{P}_Z$  the positive cone of the hyperbolic lattice  $S_Z$  containing an ample class. Note that, if  $Y$  is an Enriques surface, then  $S_Y$  is isomorphic to  $L_{10}$ .

An RDP-Enriques surface is a pair  $(Y, \rho)$  of an Enriques surface  $Y$  and a birational morphism  $\rho: Y \rightarrow \bar{Y}$  to a surface  $\bar{Y}$  that has only rational double points as its singularities. Let  $(Y, \rho)$  be an RDP-Enriques surface, and let  $C_1, \dots, C_n$  be the smooth rational curves contracted by  $\rho$ . We denote by  $\Phi_\rho \subset S_Y$  the set of classes of  $C_1, \dots, C_n$ . Then  $\Phi_\rho$  is an ADE-configuration of roots of  $S_Y$ , and  $\tau(\Phi_\rho)$  is the ADE-type of the rational double points on  $\bar{Y}$ .

**Definition 1.1.** Two RDP-Enriques surfaces  $(Y, \rho)$  and  $(Y', \rho')$  are said to be *equivalent* if there exists an isometry  $S_Y \xrightarrow{\sim} S_{Y'}$  of lattices that maps  $\mathcal{P}_Y$  to  $\mathcal{P}_{Y'}$  and  $\Phi_\rho$  to  $\Phi_{\rho'}$  bijectively.

In this paper, we classify all equivalence classes of RDP-Enriques surfaces  $(Y, \rho)$ . The ADE-type  $\tau(\Phi_\rho)$  is a principal invariant of the equivalence class, but this invariant is not enough to distinguish all the equivalence classes.

Our first main theorem is the following purely lattice-theoretic result. Let  $\Phi_f$  be an ADE-configuration of roots in  $L_{10}$ . We denote by  $R_f$  the root sublattice of  $L_{10}$  generated by  $\Phi_f$ , and by  $\bar{R}_f$  the primitive closure of  $R_f$  in  $L_{10}$ .

**Definition 1.2.** Two ADE-configurations of roots  $\Phi_f \subset L_{10}$  and  $\Phi_{f'} \subset L_{10}$  are said to be *equivalent* if there exists an isometry  $g \in O^+(L_{10})$  that maps  $\Phi_f$  to  $\Phi_{f'}$  bijectively.

**Theorem 1.3.** (1) For any ADE-configuration  $\Phi_f$  of roots in  $L_{10}$ , the lattice  $\bar{R}_f$  is a root lattice. Two ADE-configurations of roots  $\Phi_f \subset L_{10}$  and  $\Phi_{f'} \subset L_{10}$  are equivalent if and only if  $\tau(\Phi_f) = \tau(\Phi_{f'})$  and  $\tau(\bar{R}_f) = \tau(\bar{R}_{f'})$ .

(2) Let  $(t, \bar{t})$  be a pair of ADE-types. Then there exists an ADE-configuration of roots  $\Phi_f \subset L_{10}$  such that  $(\tau(\Phi_f), \tau(\bar{R}_f)) = (t, \bar{t})$  if and only if the following hold;

- (i)  $R(t)$  is of rank  $< 10$ ,
- (ii)  $R(\bar{t})$  is an even overlattice of  $R(t)$ , and
- (iii) the Dynkin diagram of  $\bar{t}$  is a sub-diagram of the Dynkin diagram of  $E_{10}$ .

There exist exactly 184 equivalence classes of ADE-configurations of roots in  $L_{10}$ . They are given in Table 1.1. In the case where  $R_f = \bar{R}_f$ , the item  $\tau(\bar{R}_f)$  is simply denoted by  $\Phi$ .

**Corollary 1.4.** Let  $(Y, \rho)$  be an RDP-Enriques surface, and  $R_\rho$  the sublattice of  $S_Y$  generated by the set  $\Phi_\rho$  of classes of smooth rational curves contracted by  $\rho$ . Then

the primitive closure  $\overline{R}_\rho$  of  $R_\rho$  in  $S_Y$  is a root lattice. Two RDP-Enriques surfaces  $(Y, \rho)$  and  $(Y', \rho')$  are equivalent if and only if  $\tau(\Phi_\rho) = \tau(\Phi_{\rho'})$  and  $\tau(\overline{R}_\rho) = \tau(\overline{R}_{\rho'})$ .

Our next problem is to determine all equivalence classes of ADE-configurations  $\Phi_f \subset L_{10}$  that can be realized as the ADE-configuration  $\Phi_\rho \subset S_Y \cong L_{10}$  associated with an RDP-Enriques surface  $(Y, \rho)$ . Let  $(Y, \rho)$  be an RDP-Enriques surface, and let  $C_1, \dots, C_n$  be the smooth rational curves on  $Y$  contracted by  $\rho$ , so that  $\Phi_\rho = \{[C_1], \dots, [C_n]\}$ . Let  $\pi: X \rightarrow Y$  denote the universal covering of  $Y$ , and  $\varepsilon: X \rightarrow X$  the deck-transformation of the double covering  $\pi$ . Then  $\pi^*: S_Y \rightarrow S_X$  is injective, and the image  $\pi^*S_Y$  is equal to the invariant sublattice in  $S_X$  of the action of  $\varepsilon$  in  $S_X$ . The pull-back of  $C_i$  by  $\pi$  splits into the disjoint union of two smooth rational curves  $C'_i$  and  $C''_i$  on  $X$ . We put

$$\Phi_\rho^\sim := \{[C'_1], [C''_1], \dots, [C'_n], [C''_n]\} \subset S_X,$$

that is,  $\Phi_\rho^\sim$  is the set of classes of smooth rational curves on  $X$  contracted by  $\rho \circ \pi: X \rightarrow \overline{Y}$ . We denote by  $M_\rho$  the sublattice of  $S_X$  generated by  $\pi^*S_Y$  and  $\Phi_\rho^\sim$ . Then the rank of  $M_\rho$  is equal to  $10 + n$ . We then denote by  $\overline{M}_\rho$  the primitive closure of  $M_\rho$  in  $S_X$ . Note that the action of  $\varepsilon$  on  $S_X$  preserves the sublattices  $M_\rho$  and  $\overline{M}_\rho$ . We put

$$Q_{(Y, \rho)} := \overline{M}_\rho / M_\rho.$$

**Definition 1.5.** Let  $(Y', \rho')$  be another RDP-Enriques surface with the universal covering  $\pi': X' \rightarrow Y'$ . We say that  $(Y, \rho)$  and  $(Y', \rho')$  are *strongly equivalent* if there exists an isometry

$$\mu: \overline{M}_\rho \xrightarrow{\sim} \overline{M}_{\rho'}$$

with the following properties; the isometry  $\mu$  maps  $\pi^*S_Y$  to  $\pi'^*S_{Y'}$  isomorphically, and the isometry  $\mu_Y: S_Y \xrightarrow{\sim} S_{Y'}$  induced by  $\mu$  maps  $\mathcal{P}_Y$  to  $\mathcal{P}_{Y'}$  and  $\Phi_\rho$  to  $\Phi_{\rho'}$  bijectively. An isometry  $\mu: \overline{M}_\rho \xrightarrow{\sim} \overline{M}_{\rho'}$  satisfying these conditions is called a *strong-equivalence isometry*.

It is obvious that, if  $(Y, \rho)$  and  $(Y', \rho')$  are strongly equivalent, then they are equivalent. It is also obvious that a strong-equivalence isometry  $\overline{M}_\rho \xrightarrow{\sim} \overline{M}_{\rho'}$  is compatible with the actions of the Enriques involutions on  $\overline{M}_\rho$  and on  $\overline{M}_{\rho'}$ . The following lemma is proved in Section 4.2.

**Lemma 1.6.** *Let  $\mu: \overline{M}_\rho \xrightarrow{\sim} \overline{M}_{\rho'}$  be a strong-equivalence isometry. Then  $\mu$  maps  $\Phi_\rho^\sim$  to  $\Phi_{\rho'}^\sim$  bijectively. In particular, if  $(Y, \rho)$  and  $(Y', \rho')$  are strongly equivalent, then we have  $Q_{(Y, \rho)} \cong Q_{(Y', \rho')}$ .*

Our second main result is as follows.

**Theorem 1.7.** *There exist exactly 265 strong equivalence classes of RDP-Enriques surfaces  $(Y, \rho)$  with  $\Phi_\rho \neq \emptyset$ . The invariants  $\tau(\Phi_\rho)$ ,  $\tau(\overline{R}_\rho)$ , and  $Q_{(Y, \rho)}$  are given in Table 1.1.*

In Table 1.1, the group  $Q_{(Y, \rho)}$  is written in the following abbreviations:

$$0 = \{0\}, \quad n = \mathbb{Z}/n\mathbb{Z} \quad (n \leq 6), \quad 22 = (\mathbb{Z}/2\mathbb{Z})^2, \quad 222 = (\mathbb{Z}/2\mathbb{Z})^3, \quad 42 = \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}.$$

When the fourth column  $Q_{(Y, \rho)}$  is empty (—), there exist no RDP-Enriques surfaces  $(Y, \rho)$  such that  $\tau(\Phi_\rho)$  and  $\tau(\overline{R}_\rho)$  are given in the second and the third columns.

**Corollary 1.8.** *There exist exactly 175 equivalence classes of RDP-Enriques surfaces  $(Y, \rho)$  with  $\Phi_\rho \neq \emptyset$ .*

No.	$\tau(\Phi_f)$	$\tau(\overline{R}_f)$	$Q_{(Y,\rho)}$	No.	$\tau(\Phi_f)$	$\tau(\overline{R}_f)$	$Q_{(Y,\rho)}$
1	$A_1$	$\Phi$	0	51	$4A_1 + A_3$	$2A_1 + D_5$	0, 2
2	$2A_1$	$\Phi$	0	52	$4A_1 + A_3$	$A_3 + D_4$	0, 2
3	$A_2$	$\Phi$	0	53	$4A_1 + A_3$	$D_7$	2, 2, 2, 22
4	$3A_1$	$\Phi$	0	54	$3A_1 + 2A_2$	$\Phi$	0
5	$A_1 + A_2$	$\Phi$	0	55	$3A_1 + A_4$	$\Phi$	0
6	$A_3$	$\Phi$	0	56	$3A_1 + D_4$	$A_1 + D_6$	0, 2
7	$4A_1$	$\Phi$	0	57	$3A_1 + D_4$	$E_7$	2, 22
8	$4A_1$	$D_4$	0, 2	58	$2A_1 + A_2 + A_3$	$\Phi$	0
9	$2A_1 + A_2$	$\Phi$	0	59	$2A_1 + A_2 + A_3$	$A_2 + D_5$	0, 2
10	$A_1 + A_3$	$\Phi$	0	60	$2A_1 + A_5$	$\Phi$	0
11	$2A_2$	$\Phi$	0	61	$2A_1 + A_5$	$A_1 + E_6$	0, 2
12	$A_4$	$\Phi$	0	62	$2A_1 + D_5$	$\Phi$	0
13	$D_4$	$\Phi$	0	63	$2A_1 + D_5$	$D_7$	0, 2
14	$5A_1$	$\Phi$	0	64	$A_1 + 3A_2$	$\Phi$	0, 3
15	$5A_1$	$A_1 + D_4$	0, 2	65	$A_1 + 3A_2$	$A_1 + E_6$	0
16	$3A_1 + A_2$	$\Phi$	0	66	$A_1 + A_2 + A_4$	$\Phi$	0
17	$2A_1 + A_3$	$\Phi$	0	67	$A_1 + A_2 + D_4$	$\Phi$	0
18	$2A_1 + A_3$	$D_5$	0, 2	68	$A_1 + 2A_3$	$\Phi$	0
19	$A_1 + 2A_2$	$\Phi$	0	69	$A_1 + 2A_3$	$A_1 + D_6$	0, 2, 2, 4
20	$A_1 + A_4$	$\Phi$	0	70	$A_1 + 2A_3$	$E_7$	0, 2
21	$A_1 + D_4$	$\Phi$	0	71	$A_1 + A_6$	$\Phi$	0
22	$A_2 + A_3$	$\Phi$	0	72	$A_1 + D_6$	$\Phi$	0
23	$A_5$	$\Phi$	0	73	$A_1 + D_6$	$E_7$	0, 2
24	$D_5$	$\Phi$	0	74	$A_1 + E_6$	$\Phi$	0
25	$6A_1$	$2A_1 + D_4$	0, 2	75	$2A_2 + A_3$	$\Phi$	0
26	$6A_1$	$D_6$	2, 22	76	$A_2 + A_5$	$\Phi$	0, 3
27	$4A_1 + A_2$	$\Phi$	0	77	$A_2 + A_5$	$E_7$	0
28	$4A_1 + A_2$	$A_2 + D_4$	0, 2	78	$A_2 + D_5$	$\Phi$	0
29	$3A_1 + A_3$	$\Phi$	0	79	$A_3 + A_4$	$\Phi$	0
30	$3A_1 + A_3$	$A_1 + D_5$	0, 2	80	$A_3 + D_4$	$\Phi$	0
31	$2A_1 + 2A_2$	$\Phi$	0	81	$A_3 + D_4$	$D_7$	0, 2, 2
32	$2A_1 + A_4$	$\Phi$	0	82	$A_7$	$\Phi$	0
33	$2A_1 + D_4$	$\Phi$	0	83	$A_7$	$E_7$	0, 2, 2
34	$2A_1 + D_4$	$D_6$	0, 2	84	$D_7$	$\Phi$	0
35	$A_1 + A_2 + A_3$	$\Phi$	0	85	$E_7$	$\Phi$	0
36	$A_1 + A_5$	$\Phi$	0	86	$8A_1$	$A_1 + E_7$	–
37	$A_1 + A_5$	$E_6$	0, 2	87	$8A_1$	$D_8$	22
38	$A_1 + D_5$	$\Phi$	0	88	$8A_1$	$E_8$	222
39	$3A_2$	$\Phi$	0, 3	89	$6A_1 + A_2$	$A_2 + D_6$	–
40	$3A_2$	$E_6$	0	90	$5A_1 + A_3$	$A_1 + D_7$	2
41	$A_2 + A_4$	$\Phi$	0	91	$4A_1 + A_4$	$A_4 + D_4$	0
42	$A_2 + D_4$	$\Phi$	0	92	$4A_1 + D_4$	$A_1 + E_7$	2
43	$2A_3$	$\Phi$	0	93	$4A_1 + D_4$	$D_8$	2, 2
44	$2A_3$	$D_6$	0, 2, 2	94	$4A_1 + D_4$	$E_8$	22, 22
45	$A_6$	$\Phi$	0	95	$3A_1 + A_2 + A_3$	$A_1 + A_2 + D_5$	0
46	$D_6$	$\Phi$	0	96	$3A_1 + A_5$	$2A_1 + E_6$	0, 2
47	$E_6$	$\Phi$	0	97	$3A_1 + D_5$	$A_1 + D_7$	0
48	$7A_1$	$A_1 + D_6$	2	98	$2A_1 + 3A_2$	$2A_1 + E_6$	0
49	$7A_1$	$E_7$	22	99	$2A_1 + A_2 + A_4$	$\Phi$	0
50	$5A_1 + A_2$	$A_1 + A_2 + D_4$	0	100	$2A_1 + A_2 + D_4$	$A_2 + D_6$	0

TABLE 1.1. ADE-configurations of roots in  $L_{10}$  (continues)

No.	$\tau(\Phi_f)$	$\tau(\overline{R}_f)$	$Q_{(Y,\rho)}$	No.	$\tau(\Phi_f)$	$\tau(\overline{R}_f)$	$Q_{(Y,\rho)}$
101	$2A_1 + 2A_3$	$A_1 + E_7$	0, 2, 4	143	$D_8$	$\Phi$	0
102	$2A_1 + 2A_3$	$A_3 + D_5$	0, 2	144	$D_8$	$E_8$	0, 2, 2
103	$2A_1 + 2A_3$	$D_8$	2, 2, 2, 22, 22, 4, 42	145	$E_8$	$\Phi$	0
104	$2A_1 + 2A_3$	$E_8$	2, 2, 2, 22	146	$9A_1$	$A_1 + E_8$	—
105	$2A_1 + A_6$	$\Phi$	0	147	$7A_1 + A_2$	$A_2 + E_7$	—
106	$2A_1 + D_6$	$A_1 + E_7$	0, 2	148	$6A_1 + A_3$	$D_9$	—
107	$2A_1 + D_6$	$D_8$	0, 2	149	$5A_1 + D_4$	$A_1 + E_8$	—
108	$2A_1 + D_6$	$E_8$	2, 2, 2, 22	150	$4A_1 + D_5$	$D_9$	—
109	$2A_1 + E_6$	$\Phi$	0	151	$3A_1 + A_2 + D_4$	$A_2 + E_7$	—
110	$A_1 + 2A_2 + A_3$	$\Phi$	0	152	$3A_1 + 2A_3$	$A_1 + E_8$	2
111	$A_1 + A_2 + A_5$	$\Phi$	0, 3	153	$3A_1 + D_6$	$A_1 + E_8$	2
112	$A_1 + A_2 + A_5$	$A_1 + E_7$	0	154	$2A_1 + A_2 + A_5$	$A_1 + E_8$	0
113	$A_1 + A_2 + A_5$	$A_2 + E_6$	0, 2, 3, 6	155	$2A_1 + A_3 + A_4$	$A_4 + D_5$	0
114	$A_1 + A_2 + A_5$	$E_8$	0, 2	156	$2A_1 + A_3 + D_4$	$D_9$	2, 2
115	$A_1 + A_2 + D_5$	$\Phi$	0	157	$2A_1 + A_7$	$A_1 + E_8$	0, 2, 4
116	$A_1 + A_3 + A_4$	$\Phi$	0	158	$2A_1 + D_7$	$D_9$	0
117	$A_1 + A_3 + D_4$	$A_1 + D_7$	0, 2	159	$2A_1 + E_7$	$A_1 + E_8$	0
118	$A_1 + A_7$	$\Phi$	0	160	$A_1 + 4A_2$	$A_1 + E_8$	—
119	$A_1 + A_7$	$A_1 + E_7$	0, 2, 2, 4, 4	161	$A_1 + A_2 + 2A_3$	$A_2 + E_7$	0
120	$A_1 + A_7$	$E_8$	0, 2, 2	162	$A_1 + A_2 + A_6$	$\Phi$	0
121	$A_1 + D_7$	$\Phi$	0	163	$A_1 + A_2 + D_6$	$A_2 + E_7$	0
122	$A_1 + E_7$	$\Phi$	0	164	$A_1 + A_2 + E_6$	$A_1 + E_8$	0
123	$A_1 + E_7$	$E_8$	0, 2	165	$A_1 + A_3 + A_5$	$A_3 + E_6$	0, 2
124	$4A_2$	$A_2 + E_6$	0, 3	166	$A_1 + A_3 + D_5$	$A_1 + E_8$	0
125	$4A_2$	$E_8$	0	167	$A_1 + 2A_4$	$A_1 + E_8$	0
126	$2A_2 + A_4$	$\Phi$	0	168	$A_1 + A_8$	$\Phi$	0, 3
127	$A_2 + 2A_3$	$A_2 + D_6$	0, 2	169	$A_1 + A_8$	$A_1 + E_8$	0
128	$A_2 + A_6$	$\Phi$	0	170	$A_1 + 2D_4$	$A_1 + E_8$	2
129	$A_2 + D_6$	$\Phi$	0	171	$A_1 + D_8$	$A_1 + E_8$	0, 2
130	$A_2 + E_6$	$\Phi$	0, 3	172	$A_1 + E_8$	$\Phi$	0
131	$A_2 + E_6$	$E_8$	0	173	$3A_2 + A_3$	$A_3 + E_6$	0
132	$A_3 + A_5$	$\Phi$	0	174	$2A_2 + A_5$	$A_2 + E_7$	0, 3
133	$A_3 + D_5$	$\Phi$	0	175	$A_2 + A_7$	$A_2 + E_7$	0, 2
134	$A_3 + D_5$	$D_8$	0, 2, 2, 4	176	$A_2 + E_7$	$\Phi$	0
135	$A_3 + D_5$	$E_8$	0, 2	177	$3A_3$	$D_9$	2, 22
136	$2A_4$	$\Phi$	0, 5	178	$A_3 + D_6$	$D_9$	0, 2
137	$2A_4$	$E_8$	0	179	$A_3 + E_6$	$\Phi$	0
138	$A_4 + D_4$	$\Phi$	0	180	$A_4 + A_5$	$\Phi$	0
139	$A_8$	$\Phi$	0, 3	181	$A_4 + D_5$	$\Phi$	0
140	$A_8$	$E_8$	0	182	$A_9$	$\Phi$	0
141	$2D_4$	$D_8$	0, 2	183	$D_4 + D_5$	$D_9$	0
142	$2D_4$	$E_8$	2, 2, 22	184	$D_9$	$\Phi$	0

TABLE 1.1. ADE-configurations of roots in  $L_{10}$  (continued)

**Example 1.9.** There exist no RDP-Enriques surfaces  $(Y, \rho)$  with singularities of type  $\tau(\Phi_\rho) = 6A_1 + A_2$ .

**Example 1.10.** RDP-Enriques surfaces  $(Y, \rho)$  with singularities of type  $\tau(\Phi_\rho) = 2A_1 + 2A_3$  are divided into 4 equivalence classes with

$$\tau(\overline{R}_\rho) = A_1 + E_7, \quad A_3 + D_5, \quad D_8, \quad E_8,$$

and into  $3 + 2 + 7 + 4 = 16$  strong equivalence classes.

In [12], Keum and Zhang studied configurations  $\mathcal{A}$  of smooth rational curves on an Enriques surface  $Y$  with ADE-type  $cA_{p-1}$ , where  $p$  is a prime, and investigated the topological fundamental group  $\pi_1(Y \setminus \mathcal{A})$ . In [17], Rams and Schütt studied the case of  $4A_2$ , and corrected the result of [12]. The relation of the isomorphism classes of  $\pi_1(Y \setminus \mathcal{A})$  and our notion of strong equivalence relation is not clear. We observe, however, the following fact by comparing our Table 1.1 with their results (Theorem 2 and Table 2 of [12] and Theorem 4.3 of [17]): Except for the case of the  $8A_1$  (and possibly for the case of the  $7A_1$ ), the number of the isomorphism classes of  $\pi_1(Y \setminus \mathcal{A})$  and the number of strong equivalence classes coincide. For example, consider the case of  $7A_1$ . The number of the isomorphism classes of  $\pi_1(Y \setminus \mathcal{A})$  is two or three (the authors of [12] did not determine the realizability of the case  $\pi_1(Y \setminus \mathcal{A}) \cong (\mathbb{Z}/2\mathbb{Z})^4$ ), whereas we have two strong equivalence classes (Nos. 48 and 49). Therefore we guess that the case  $\pi_1(Y \setminus \mathcal{A}) \cong (\mathbb{Z}/2\mathbb{Z})^4$  is not realizable. For another example, consider the case of  $4A_2$ . Theorem 4.3 of [17] says that there exist three isomorphism classes of  $\pi_1(Y \setminus \mathcal{A})$ , two of which are realized by the same Enriques surface. We also have  $3 = 2 + 1$  strong equivalence classes (Nos. 124 and 125). For the case of  $8A_1$ , Table 2 of [12] indicates only one isomorphism class of  $\pi_1(Y \setminus \mathcal{A})$ , whereas we have two strong equivalence classes (Nos. 87 and 88).

An important class of involutions of  $K3$  surfaces other than Enriques involutions comes from the sextic double plane models. The classification of ADE-types of rational double points on normal sextic double planes was given by Yang [27]. A finer classification of rational double points on normal sextic double planes was given in [20] in the relation with the topology of Zariski pairs of plane curves (see [2], [21]). This classification was further refined to the complete description of connected components of the equisingular families of irreducible sextic plane curves with only simple singularities by Akyol and Degtyarev [1]. The present article may be regarded as the Enriques counterpart of these studies of plane sextic curves.

This paper is organized as follows. In Section 2, we review preliminary results about lattices. Some algorithms for negative-definite root lattices are presented, and the notion of chambers in a hyperbolic lattice is introduced. In Section 3, we study the lattice  $L_{10}$  in detail, and prove Theorem 1.3 by the generalized Borcherds algorithm ([5], [6], [22]). The classical result due to Vinberg [26] plays an important role. In Section 4, we give a method to enumerate all the strong equivalence classes of RDP-Enriques surfaces, explain how to carry out this method, and prove Theorem 1.7.

For the computation, we used GAP [9]. A computational data is available from the author's webpage [25]. In particular, the isomorphism class of the lattice  $\overline{M}_\rho$  for each strongly equivalence class of RDP-Enriques surfaces is given explicitly in [25]. Since we have  $\overline{M}_\rho = S_X$  when  $(Y, \rho)$  corresponds to a general point of an irreducible component of the moduli of RDP-Enriques surfaces with a fixed ADE-type, this data will be useful in the study of the moduli of RDP-Enriques surfaces.

The author thanks Professors Matthias Schütt and Hisanori Ohashi for many discussions. Thanks are also due to the referees for many helpful comments on the first version of the manuscript.

## 2. PRELIMINARIES

**2.1. An ADE-configuration.** A *Dynkin configuration* is a finite set  $\Phi$  with a map

$$\langle \cdot, \cdot \rangle: \Phi \times \Phi \rightarrow \{-2, 0, 1\}$$

such that  $\langle x, y \rangle = \langle y, x \rangle$  for all  $x, y \in \Phi$ ,  $\langle x, x \rangle = -2$  for all  $x \in \Phi$ , and  $\langle x, y \rangle \in \{0, 1\}$  for all  $x, y \in \Phi$  with  $x \neq y$ . With a Dynkin configuration  $\Phi = \{r_1, \dots, r_n\}$ , we associate its *Dynkin diagram*, which is a graph whose set of vertices is  $\Phi$  and whose set of edges is the set of pairs  $\{r_i, r_j\}$  such that  $\langle r_i, r_j \rangle = 1$ . We say that a Dynkin configuration is an *ADE-configuration* if every connected component of its Dynkin diagram is of type  $A_l$  ( $l \geq 1$ ),  $D_m$  ( $m \geq 4$ ), or  $E_n$  ( $n = 6, 7, 8$ ). We define the *ADE-type*  $\tau(\Phi)$  of an ADE-configuration  $\Phi$  to be the sum of the types of the connected components of its Dynkin diagram. Let  $\Phi$  and  $\Phi'$  be Dynkin configurations. An isomorphism from  $\Phi$  to  $\Phi'$  is a bijection  $\Phi \xrightarrow{\sim} \Phi'$  that preserves  $\langle \cdot, \cdot \rangle$ . The isomorphism class of an ADE-configuration is uniquely determined by its ADE-type. We denote by  $\text{Aut}(\Phi)$  the group of automorphisms of  $\Phi$ , which we let act on  $\Phi$  from the *right*.

**2.2. A lattice.** Let  $L$  be a free  $\mathbb{Z}$ -module of finite rank, and  $R$  a submodule of  $L$ . The *primitive closure*  $\bar{R}$  of  $R$  in  $L$  is the intersection of  $R \otimes \mathbb{Q}$  and  $L$  in  $L \otimes \mathbb{Q}$ . We say that  $R$  is *primitive in  $L$*  if  $R = \bar{R}$ .

A *lattice* is a free  $\mathbb{Z}$ -module  $L$  of finite rank with a non-degenerate symmetric bilinear form

$$\langle \cdot, \cdot \rangle: L \times L \rightarrow \mathbb{Z}.$$

Let  $L$  be a lattice. We say that  $L$  is *even* if  $\langle x, x \rangle \in 2\mathbb{Z}$  for all  $x \in L$ . The group of isometries of  $L$  is denoted by  $O(L)$ . We let  $O(L)$  act on  $L$  from the *right*. An *embedding* of a Dynkin configuration  $\Phi$  into  $L$  is an injection  $\Phi \hookrightarrow L$  that preserves  $\langle \cdot, \cdot \rangle$ . We define the *dual lattice*  $L^\vee$  of  $L$  by

$$L^\vee := \{ v \in L \otimes \mathbb{Q} \mid \langle x, v \rangle \in \mathbb{Z} \text{ for all } x \in L \}.$$

The finite abelian group  $A_L := L^\vee/L$  is called the *discriminant group* of  $L$ . The group  $O(L)$  acts on  $A_L$  from the right. We say that  $L$  is *unimodular* if  $A_L$  is trivial. The *signature* of  $L$  is the signature of the real quadratic space  $L \otimes \mathbb{R}$ . Suppose that  $L$  is of rank  $n > 0$ . We say that  $L$  is *hyperbolic* if the signature is  $(1, n - 1)$ , and is *negative-definite* if the signature is  $(0, n)$ .

**2.3. Roots and reflections.** Let  $L$  be an even lattice. A *root* of  $L$  is a vector  $r \in L$  such that  $\langle r, r \rangle = -2$ . The set of roots of  $L$  is denoted by  $\text{Roots}(L)$ . A root  $r$  of  $L$  defines an isometry

$$s_r: x \mapsto x + \langle x, r \rangle r$$

of  $L$ , which is called the *reflection* associated with  $r$ . We denote by  $W(L)$  the subgroup of  $O(L)$  generated by all the reflections associated with the roots, and call it the *Weyl group* of  $L$ . We say that  $L$  is a *root lattice* if  $L$  is generated by roots. Let  $\Phi$  be a subset of  $\text{Roots}(L)$ . We denote by  $W(\Phi, L)$  the subgroup of  $W(L)$  generated by all the reflections  $s_r$  associated with  $r \in \Phi$ .

**2.4. A negative-definite root lattice.** Let  $\Phi$  be an ADE-configuration. Extending  $\langle \cdot, \cdot \rangle: \Phi \times \Phi \rightarrow \mathbb{Z}$  by linearity to the bilinear form on the free  $\mathbb{Z}$ -module generated by  $\Phi$ , we obtain a negative-definite root lattice of rank  $|\Phi|$ , which we will denote by  $\langle \Phi \rangle$ . Conversely, let  $R$  be a negative-definite root lattice. Then  $R$  has a basis  $\Phi_R$  consisting of roots that form an ADE-configuration, which we call an ADE-basis of  $R$ . We define the ADE-type  $\tau(R)$  of  $R$  to be the ADE-type  $\tau(\Phi_R)$  of an ADE-basis  $\Phi_R$  of  $R$ . In the following, we describe the set of all ADE-bases of a negative-definite root lattice. See [8, Chapter 1] or [10, Chapter 1] for the proof.

Let  $R$  be a negative-definite root lattice. For a root  $r$  of  $R$ , we denote by  $r^\perp$  the hyperplane of  $R \otimes \mathbb{R}$  defined by  $\langle x, r \rangle = 0$ . We then put

$$(R \otimes \mathbb{R})^\circ := (R \otimes \mathbb{R}) \setminus \bigcup r^\perp,$$

where  $r$  runs through the finite set  $\text{Roots}(R)$ . Let  $\Gamma$  be a connected component of  $(R \otimes \mathbb{R})^\circ$ , and  $\bar{\Gamma}$  the closure of  $\Gamma$  in  $R \otimes \mathbb{R}$ . Then the set  $\Phi_\Gamma$  of all  $r \in \text{Roots}(R)$  such that  $\langle x, r \rangle > 0$  for any  $x \in \Gamma$  and that  $r^\perp \cap \bar{\Gamma}$  contains a non-empty open subset of  $r^\perp$  form an ADE-basis of  $R$ , and the mapping  $\Gamma \mapsto \Phi_\Gamma$  gives a bijection from the set of connected components of  $(R \otimes \mathbb{R})^\circ$  to the set of ADE-bases of  $R$ . For  $x \in (R \otimes \mathbb{R})^\circ$ , we denote by  $\Gamma(x)$  the connected component of  $(R \otimes \mathbb{R})^\circ$  containing  $x$ . Let  $\Phi = \{r_1, \dots, r_n\}$  be an ADE-basis of  $R$ . We put

$$c := r_1^\vee + \dots + r_n^\vee \in (R \otimes \mathbb{R})^\circ,$$

where  $r_1^\vee, \dots, r_n^\vee$  are the basis of  $R^\vee$  dual to the basis  $r_1, \dots, r_n$  of  $R$ . Then the connected component of  $(R \otimes \mathbb{R})^\circ$  corresponding to the ADE-basis  $\Phi$  is  $\Gamma(c)$ . It is obvious that we have a natural embedding  $\text{Aut}(\Phi) \hookrightarrow \text{O}(R)$  whose image is

$$\text{Stab}(\Gamma(c), R) := \{g \in \text{O}(R) \mid \Gamma(c)^g = \Gamma(c)\} = \{g \in \text{O}(R) \mid c^g = c\}.$$

Since  $W(R)$  acts on the set of connected components of  $(R \otimes \mathbb{R})^\circ$  simple-transitively, we have a *splitting* exact sequence

$$(2.1) \quad 1 \rightarrow W(R) \rightarrow \text{O}(R) \xrightarrow{\kappa} \text{Aut}(\Phi) \rightarrow 1.$$

**Algorithm 2.1.** Let  $u$  and  $v$  be points of  $(R \otimes \mathbb{Q}) \cap (R \otimes \mathbb{R})^\circ$ . This algorithm finds the unique element  $g \in W(R)$  that maps  $\Gamma(u)$  to  $\Gamma(v)$ . Let  $\xi$  be a sufficiently general element of  $R \otimes \mathbb{Q}$ , and let  $\varepsilon$  be a sufficiently small positive rational number. We consider the open line segment in  $R \otimes \mathbb{R}$  drawn by the point

$$p(t) := u + t(v + \varepsilon\xi),$$

where  $t$  moves in the set of positive real numbers. Let  $\{r_1, \dots, r_N\}$  be the set of roots  $r_i$  of  $R$  such that  $\langle u, r_i \rangle < 0$  and  $\langle v, r_i \rangle > 0$ . For each  $r_i$ , let  $t_i$  be the unique rational number such that  $\langle p(t_i), r_i \rangle = 0$ . Since the perturbation vector  $\xi$  is general, we can assume that  $t_1, \dots, t_N$  are distinct. We sort  $r_1, \dots, r_N$  in such a way that  $t_1 < \dots < t_N$ . Then  $g := s_{r_1} \dots s_{r_N} \in W(R)$  satisfies  $\Gamma(u)^g = \Gamma(v)$ .  $\blacksquare$

As applications of Algorithm 2.1, we obtain the following algorithms. As noted above, the stabilizer subgroup  $\text{Stab}(\Gamma(c), R)$  is canonically identified with  $\text{Aut}(\Phi)$ .

**Algorithm 2.2.** Let an isometry  $g \in \text{O}(R)$  be given. The image  $\kappa(g) \in \text{Aut}(\Phi)$  of  $g$  by the homomorphism  $\kappa$  in (2.1) is calculated by applying Algorithm 2.1 to  $u = c^g$  and  $v = c$ . We find  $h \in W(R)$  such that  $\Gamma(c)^{gh} = \Gamma(c)$ . Hence we have  $\kappa(g) = gh$ .  $\blacksquare$



**Algorithm 2.3.** Applying Algorithm 2.1 to  $u = c$  and  $v = -c$ , we find the element  $l \in W(R)$  such that  $\Gamma(c)^l = -\Gamma(c)$ . This element  $l$  is the *longest element* of the Coxeter group  $W(R)$ . (See [10, Section 1.8].)  $\blacksquare$

Another method to obtain an ADE-basis of  $R$  is as follows. We put

$$\mathrm{Hom}(R, \mathbb{R})^\circ := \{ \ell \in \mathrm{Hom}(R, \mathbb{R}) \mid \ell(r) \neq 0 \text{ for any } r \in \mathrm{Roots}(R) \},$$

and for  $\ell \in \mathrm{Hom}(R, \mathbb{R})^\circ$ , we put  $\mathrm{Roots}(R)_{\ell > 0} := \{ r \in \mathrm{Roots}(R) \mid \ell(r) > 0 \}$ .

**Definition 2.4.** Let  $S$  be a subset of  $\mathrm{Roots}(R)_{\ell > 0}$ . We say that  $r \in S$  is *indecomposable in  $S$*  if  $r$  is not written as a linear combination  $\sum a_i r_i$  of elements  $r_i \in S$  with  $a_i \in \mathbb{Z}_{\geq 0}$  such that  $\sum a_i > 1$ .

Let  $\Phi_{\ell > 0}$  be the set of roots  $r \in \mathrm{Roots}(R)_{\ell > 0}$  indecomposable in  $\mathrm{Roots}(R)_{\ell > 0}$ . Then  $\Phi_{\ell > 0}$  is an ADE-basis of  $R$ , and the mapping  $\ell \mapsto \Phi_{\ell > 0}$  gives a bijection from the set of connected components of  $\mathrm{Hom}(R, \mathbb{R})^\circ$  to the set of ADE-bases of  $R$ . This correspondence  $\ell \mapsto \Phi_{\ell > 0}$  will be used in Section 4.

**2.5. An even hyperbolic lattice.** Let  $L$  be an even hyperbolic lattice. A *positive cone* is one of the two connected components of the space  $\{x \in L \otimes \mathbb{R} \mid \langle x, x \rangle > 0\}$ . Let  $\mathcal{P}$  be a positive cone. We denote by  $O^+(L)$  the stabilizer subgroup of  $\mathcal{P}$  in  $O(L)$ . For a non-zero vector  $v \in L \otimes \mathbb{R}$ , we put

$$H_v^+ := \{ x \in \mathcal{P} \mid \langle v, x \rangle \geq 0 \}, \quad (v)^\perp := \{ x \in \mathcal{P} \mid \langle v, x \rangle = 0 \}.$$

Then  $(v)^\perp \neq \emptyset$  if and only if  $\langle v, v \rangle < 0$ . Note that the Weyl group  $W(L)$  acts on  $\mathcal{P}$ . A *standard fundamental domain of the action of  $W(L)$  on  $\mathcal{P}$*  is the closure in  $\mathcal{P}$  of a connected component of

$$\mathcal{P} \setminus \bigcup (r)^\perp,$$

where  $r$  runs through  $\mathrm{Roots}(L)$ . Then  $W(L)$  acts on the set of standard fundamental domains simple-transitively. Let  $\Delta$  be a standard fundamental domain. We put  $\mathrm{Aut}(\Delta) := \{g \in O^+(L) \mid \Delta^g = \Delta\}$ . Then we have a *splitting* exact sequence

$$(2.2) \quad 1 \rightarrow W(L) \rightarrow O^+(L) \rightarrow \mathrm{Aut}(\Delta) \rightarrow 1,$$

and we have a tessellation

$$(2.3) \quad \mathcal{P} = \bigcup_{g \in W(L)} \Delta^g.$$

**2.6. Chambers.** Let  $L$  be an even hyperbolic lattice, and  $\mathcal{P}$  a positive cone of  $L$ . A closed subset  $\mathcal{C}$  of  $\mathcal{P}$  is called a *chamber* if the interior  $\mathcal{C}^\circ$  of  $\mathcal{C}$  in  $\mathcal{P}$  is non-empty and there exists a set of vectors  $\mathcal{H} \subset L \otimes \mathbb{R}$  such that the family of hyperplanes  $\{(v)^\perp \mid v \in \mathcal{H}\}$  of  $\mathcal{P}$  is locally finite in  $\mathcal{P}$  and that

$$(2.4) \quad \mathcal{C} = \bigcap_{v \in \mathcal{H}} H_v^+.$$

Let  $\mathcal{C}$  be a chamber defined by a set of vectors  $\mathcal{H} \subset L \otimes \mathbb{R}$  as in (2.4). A closed subset  $F$  of  $\mathcal{C}$  is called a *face* of  $\mathcal{C}$  if  $F \neq \emptyset$ ,  $F \cap \mathcal{C}^\circ = \emptyset$ , and there exists a subset  $\mathcal{H}_F$  of  $\mathcal{H}$  such that

$$F = \mathcal{C} \cap \bigcap_{v \in \mathcal{H}_F} (v)^\perp.$$

If  $F$  is a face, then we have a unique linear subspace  $V$  of  $L \otimes \mathbb{R}$  such that  $V \cap F$  contains a non-empty open subset of  $V$ . We say that  $V$  or  $\mathcal{P} \cap V$  the *supporting*

linear subspace of  $F$ . The *codimension* of a face is defined to be the codimension of its supporting linear subspace in  $L \otimes \mathbb{R}$  or in  $\mathcal{P}$ . A face of codimension 1 is called a *wall*. Let  $F$  be a wall of  $\mathcal{C}$ . A vector  $v \in L \otimes \mathbb{R}$  is said to *define the wall  $F$*  if  $\mathcal{C}$  is contained in  $H_v^+$  and  $F = \mathcal{C} \cap (v)^\perp$  holds.

**2.7. The discriminant form and overlattices.** Let  $L$  be an even lattice. Recall that  $A_L := L^\vee/L$ . Then the natural  $\mathbb{Q}$ -valued symmetric bilinear form on  $L^\vee$  defines a finite quadratic form

$$q_L : A_L \rightarrow \mathbb{Q}/2\mathbb{Z},$$

which we call the *discriminant form* of  $L$ . See Nikulin [14] for the basic properties of the discriminant form. We have a natural homomorphism  $O(L) \rightarrow O(q_L)$ , where  $O(q_L)$  is the automorphism group of the finite quadratic form  $q_L$ . An *even overlattice* of  $L$  is a submodule  $M$  of  $L^\vee$  containing  $L$  such that the restriction of the natural  $\mathbb{Q}$ -valued symmetric bilinear form on  $L^\vee$  makes  $M$  an even lattice. By definition, we have the following:

**Proposition 2.5.** *The map  $M \mapsto M/L$  gives a bijection from the set of even overlattices  $M$  of  $L$  to the set of totally isotropic subgroups of  $q_L$ .  $\square$*

Note that  $O(L)$  acts on the set of even overlattices of  $L$  from the right.

**Proposition 2.6.** *Suppose that the signature of  $L$  is  $(s_+, s_-)$ . Let  $H$  be an even unimodular lattice of signature  $(h_+, h_-)$ . Then  $L$  can be embedded into  $H$  primitively if and only if there exists an even lattice of signature  $(h_+ - s_+, h_- - s_-)$  whose discriminant form is isomorphic to  $-q_L$ .  $\square$*

*Remark 2.7.* The signature  $(h_+ - s_+, h_- - s_-)$  and the isomorphism class of the discriminant form  $-q_L$  determine a genus of even lattices. There exist various versions of the criterion to determine whether a genus given by signature and a finite quadratic form is empty or not. See, for example, Nikulin [14], Conway-Sloane [7, Chapter 15], Miranda-Morrison [13]. This criterion has been applied to many problems of  $K3$  surfaces. See, for example, [1, 19, 20, 23, 27].

### 3. THE LATTICE $L_{10}$

For an embedding  $f: \Phi \hookrightarrow L_{10}$  of an ADE-configuration  $\Phi$  into  $L_{10}$ , we denote by  $\Phi_f$  the image of  $\Phi$  by  $f$ . We extend Definition 1.2 to the equivalence relation of embeddings of ADE-configurations into  $L_{10}$ .

**Definition 3.1.** Let  $f: \Phi \hookrightarrow L_{10}$  and  $f': \Phi' \hookrightarrow L_{10}$  be embeddings of ADE-configurations  $\Phi$  and  $\Phi'$ . We say that  $f$  and  $f'$  are *equivalent* if there exist an isomorphism  $\gamma: \Phi \xrightarrow{\sim} \Phi'$  and an isometry  $g \in O^+(L_{10})$  that make the following diagram commutative:

$$\begin{array}{ccc} \Phi & \xrightarrow{f} & L_{10} \\ \gamma \downarrow & & \downarrow g \\ \Phi' & \xrightarrow{f'} & L_{10}. \end{array}$$

The purpose of this section is to prove Theorem 1.3.

### 3.1. Negative-definite primitive root sublattices of $L_{10}$ .

**Definition 3.2.** Let  $\mathcal{N}$  denote the set of all negative-definite primitive root sublattices of  $L_{10}$ , on which  $O^+(L_{10})$  acts from the right.

We calculate the set  $\mathcal{N}/O^+(L_{10})$  of orbits of this action. Recall that the lattice  $L_{10}$  has a basis  $E_{10} = \{e_1, \dots, e_{10}\}$  consisting of roots that form the Dynkin diagram in Figure 1.1. We fix, once and for all, the positive cone  $\mathcal{P}_{10}$  that contains the vector

$$c_0 := e_1^\vee + \dots + e_{10}^\vee$$

of square-norm 1240, where  $e_1^\vee, \dots, e_{10}^\vee$  are the basis of  $L_{10}^\vee = L_{10}$  dual to the basis  $e_1, \dots, e_{10}$ . For simplicity, we put, for a subset  $S$  of  $L_{10} \otimes \mathbb{R}$ ,

$$(3.1) \quad [S]^\perp := \{v \in L_{10} \mid \langle v, x \rangle = 0 \text{ for all } x \in S\},$$

$$(3.2) \quad (S)^\perp := \{v \in \mathcal{P}_{10} \mid \langle v, x \rangle = 0 \text{ for all } x \in S\}.$$

We consider the chamber

$$\Delta_0 := \{x \in \mathcal{P}_{10} \mid \langle x, e_i \rangle \geq 0 \text{ for all } i = 1, \dots, 10\},$$

which contains  $c_0$  in its interior. Vinberg [26] proved the following:

**Theorem 3.3** (Vinberg). *Each  $e_i$  defines a wall of  $\Delta_0$ . The chamber  $\Delta_0$  is a standard fundamental domain of the action of  $W(L_{10})$  on  $\mathcal{P}_{10}$ .  $\square$*

**Definition 3.4.** We call a standard fundamental domain of the action of  $W(L_{10})$  on  $\mathcal{P}_{10}$  a *Vinberg chamber*. We denote by  $\mathcal{V}$  the set of all Vinberg chambers.

The following easy fact is used frequently in this section: Let  $r_1, \dots, r_n$  be roots of  $L_{10}$  such that the linear subspace  $\mathcal{P}' := (\{r_1, \dots, r_n\})^\perp$  of  $\mathcal{P}_{10}$  is non-empty. Let  $\Delta$  be a Vinberg chamber. If  $\mathcal{P}' \cap \Delta$  contains a non-empty open subset of  $\mathcal{P}'$ , then  $\mathcal{P}' \cap \Delta$  is a face of  $\Delta$ , and its supporting linear subspace is  $\mathcal{P}'$ .

Since the Dynkin diagram in Figure 1.1 has no symmetries, we see from (2.2) that  $O^+(L_{10})$  is equal to  $W(L_{10})$ . In particular, we have the following:

**Proposition 3.5.** *The map  $g \mapsto \Delta_0^g$  is a bijection from  $O^+(L_{10})$  to  $\mathcal{V}$ .  $\square$*

We denote by  $\gamma: \mathcal{V} \rightarrow O^+(L_{10})$  the inverse map of  $g \mapsto \Delta_0^g$ , that is,  $\gamma(\Delta)$  is the unique element of  $O^+(L_{10})$  such that

$$\Delta_0^{\gamma(\Delta)} = \Delta.$$

The following lemma is easy to prove:

**Lemma 3.6.** (1) *A subset  $\Sigma$  of  $E_{10}$  is an ADE-configuration of roots if and only if  $\Sigma \neq \emptyset$ ,  $\Sigma \neq E_{10}$ , and  $\Sigma \neq \{e_1, \dots, e_9\}$ . (2) *Let  $\overline{\mathcal{P}}_{10}$  and  $\overline{\Delta}_0$  denote the closure of  $\mathcal{P}_{10}$  and  $\Delta_0$  in  $L_{10} \otimes \mathbb{R}$ , respectively. Then  $\overline{\Delta}_0 \cap (\overline{\mathcal{P}}_{10} \setminus \mathcal{P}_{10})$  is equal to the half-line  $\mathbb{R}_{\geq 0}e_{10}^\vee = (\{e_1, \dots, e_9\})^\perp \cap \overline{\mathcal{P}}_{10}$ .  $\square$**

Let  $2^{E_{10}}$  be the power set of  $E_{10} = \{e_1, \dots, e_{10}\}$ . We consider the set

$$\mathcal{S} := 2^{E_{10}} \setminus \{\emptyset, E_{10}, \{e_1, \dots, e_9\}\}.$$

Since  $\overline{\Delta}_0$  is the cone over a 9-dimensional simplex, we see that

$$\Sigma \mapsto F_\Sigma := (\Sigma)^\perp \cap \Delta_0$$

is a bijection from  $\mathcal{S}$  to the set of faces of  $\Delta_0$ . For an element  $\Sigma$  of  $\mathcal{S}$ , we denote by  $\langle \Sigma \rangle$  the sublattice of  $L_{10}$  generated by  $\Sigma$ . Since  $\Sigma$  is a subset of the basis  $E_{10}$  of  $L_{10}$ , the sublattice  $\langle \Sigma \rangle$  is primitive in  $L_{10}$ . Therefore we obtain the following:

**Lemma 3.7.** *Let  $\Sigma$  be an element of  $\mathcal{S}$ . Then  $\langle \Sigma \rangle \in \mathcal{N}$  and  $\langle \Sigma \rangle = [F_\Sigma]^\perp$ .  $\square$*

**Lemma 3.8.** *Let  $R$  be a negative-definite root sublattice of  $L_{10}$ , and  $\bar{R}$  the primitive closure of  $R$  in  $L_{10}$ . Then there exists an isometry  $g \in \mathcal{O}^+(L_{10})$  such that  $\bar{R}^g = \langle \Sigma \rangle$  for some  $\Sigma \in \mathcal{S}$ . In particular, the lattice  $\bar{R}$  is a root lattice.*

*Proof.* Suppose that  $R$  is generated by roots  $r_1, \dots, r_n$ . Then

$$\mathcal{P}(R) := (\{r_1, \dots, r_n\})^\perp = ([R]^\perp \otimes \mathbb{R}) \cap \mathcal{P}_{10}$$

is non-empty, because  $\mathcal{P}(R)$  is a positive cone of the hyperbolic lattice  $[R]^\perp$ . Since  $\mathcal{P}(R)$  is contained in a hyperplane  $(r_1)^\perp$ , we see that  $\mathcal{P}(R)$  is disjoint from the interior of any Vinberg chamber. Since  $\mathcal{P}_{10}$  is tessellated by Vinberg chambers, there exists a Vinberg chamber  $\Delta$  such that  $\mathcal{P}(R) \cap \Delta$  contains a non-empty open subset of  $\mathcal{P}(R)$ . Therefore  $\mathcal{P}(R) \cap \Delta$  is a face of  $\Delta$ . Then  $\gamma(\Delta)^{-1} \in \mathcal{O}^+(L_{10})$  maps  $\mathcal{P}(R) \cap \Delta$  to a face  $F_\Sigma$  of  $\Delta_0$  associated with some  $\Sigma \in \mathcal{S}$ . Hence  $\gamma(\Delta)^{-1}$  maps  $\bar{R} = [\mathcal{P}(R) \cap \Delta]^\perp$  to  $\langle \Sigma \rangle = [F_\Sigma]^\perp$ .  $\square$

**Corollary 3.9.** *The map  $\Sigma \mapsto \langle \Sigma \rangle$  induces a surjection from  $\mathcal{S}$  to  $\mathcal{N}/\mathcal{O}^+(L_{10})$ .  $\square$*

For  $\Sigma \in \mathcal{S}$ , we put

$$\mathcal{P}(\Sigma) := (\Sigma)^\perp = ([\Sigma]^\perp \otimes \mathbb{R}) \cap \mathcal{P}_{10}.$$

**Definition 3.10.** For  $\Sigma, \Sigma' \in \mathcal{S}$ , we write  $\Sigma \sim \Sigma'$  if there exists an isometry  $g \in \mathcal{O}^+(L_{10})$  such that  $\langle \Sigma \rangle^g = \langle \Sigma' \rangle$ , or equivalently,  $\mathcal{P}(\Sigma)^g = \mathcal{P}(\Sigma')$ .

In the following, we fix an element  $\Sigma \in \mathcal{S}$ , and calculate the set  $\{\Sigma' \in \mathcal{S} \mid \Sigma' \sim \Sigma\}$  and a finite generating set of the stabilizer subgroup

$$\text{Stab}(\langle \Sigma \rangle, L_{10}) := \{g \in \mathcal{O}^+(L_{10}) \mid \langle \Sigma \rangle^g = \langle \Sigma \rangle\} = \{g \in \mathcal{O}^+(L_{10}) \mid \mathcal{P}(\Sigma)^g = \mathcal{P}(\Sigma)\}$$

of  $\langle \Sigma \rangle$  in  $\mathcal{O}^+(L_{10})$ . We put

$$\mathcal{V}_\Sigma := \{ \Delta \in \mathcal{V} \mid \mathcal{P}(\Sigma) \cap \Delta \text{ contains a non-empty open subset of } \mathcal{P}(\Sigma) \}.$$

**Definition 3.11.** If  $\Delta \in \mathcal{V}_\Sigma$ , then  $\mathcal{P}(\Sigma) \cap \Delta$  is a chamber in the positive cone  $\mathcal{P}(\Sigma)$  of the hyperbolic lattice  $[\Sigma]^\perp$ . A closed subset  $D$  of  $\mathcal{P}(\Sigma)$  is said to be an *induced chamber* if  $D$  is written as  $\mathcal{P}(\Sigma) \cap \Delta$  by some  $\Delta \in \mathcal{V}_\Sigma$ .

By definition, the positive cone  $\mathcal{P}(\Sigma)$  of the hyperbolic lattice  $[\Sigma]^\perp$  is covered by induced chambers in such a way that, if  $D$  and  $D'$  are distinct induced chambers, then their interiors in  $\mathcal{P}(\Sigma)$  are disjoint. Let  $\Delta$  be an arbitrary element of  $\mathcal{V}_\Sigma$ . Looking at the tessellation of  $\mathcal{P}_{10}$  by Vinberg chambers locally around an interior point of the induced chamber  $\mathcal{P}(\Sigma) \cap \Delta$ , we obtain the following:

**Lemma 3.12.** *Let  $W(\Sigma, L_{10})$  denote the subgroup of  $\mathcal{O}^+(L_{10})$  generated by the reflections  $s_r$  associated with the roots  $r \in \Sigma$ . Let  $\Delta$  be an element of  $\mathcal{V}_\Sigma$ . Then  $W(\Sigma, L_{10})$  acts on the set  $\{\Delta' \in \mathcal{V}_\Sigma \mid \mathcal{P}(\Sigma) \cap \Delta = \mathcal{P}(\Sigma) \cap \Delta'\}$  simple-transitively.  $\square$*

Note that, if  $\Delta \in \mathcal{V}_\Sigma$ , then  $\gamma(\Delta)^{-1}$  maps the induced chamber  $\mathcal{P}(\Sigma) \cap \Delta$  of  $\mathcal{P}(\Sigma)$  to a face of  $\Delta_0$ . Hence we can define a mapping  $\sigma: \mathcal{V}_\Sigma \rightarrow \mathcal{S}$  by the property

$$(3.3) \quad (\mathcal{P}(\sigma(\Delta)) \cap \Delta_0)^{\gamma(\Delta)} = \mathcal{P}(\Sigma) \cap \Delta,$$

or equivalently, by the equality  $F_{\sigma(\Delta)}^{\gamma(\Delta)} = \mathcal{P}(\Sigma) \cap \Delta$ . Taking the orthogonal complement of the both sides, we obtain

$$(3.4) \quad \langle \sigma(\Delta) \rangle^{\gamma(\Delta)} = \langle \Sigma \rangle$$

**Lemma 3.13.** *We have  $\{\Sigma' \in \mathcal{S} \mid \Sigma' \sim \Sigma\} = \{\sigma(\Delta) \mid \Delta \in \mathcal{V}_\Sigma\}$ .*

*Proof.* By (3.4), we have  $\sigma(\Delta) \sim \Sigma$  for any  $\Delta \in \mathcal{V}_\Sigma$ . Suppose that  $\Sigma' \sim \Sigma$ , and let  $h \in \mathcal{O}^+(L_{10})$  be an isometry such that  $\mathcal{P}(\Sigma')^h = \mathcal{P}(\Sigma)$ . We put  $\Delta := \Delta_0^h$  so that  $\gamma(\Delta) = h$ . Since  $(\mathcal{P}(\Sigma') \cap \Delta_0)^h = \mathcal{P}(\Sigma) \cap \Delta$  contains a non-empty open subset of  $\mathcal{P}(\Sigma)$ , we have  $\Delta \in \mathcal{V}_\Sigma$  and  $\Sigma' = \sigma(\Delta)$  by the defining property (3.3) of  $\sigma$ .  $\square$

Using Lemma 3.12, we can prove the following:

**Lemma 3.14.** *For  $\Delta, \Delta' \in \mathcal{V}_\Sigma$ , if  $\mathcal{P}(\Sigma) \cap \Delta = \mathcal{P}(\Sigma) \cap \Delta'$ , then  $\sigma(\Delta) = \sigma(\Delta')$ .  $\square$*

Let  $\Delta$  be an element of  $\mathcal{V}_\Sigma$ , and let  $D := \mathcal{P}(\Sigma) \cap \Delta$  be the corresponding induced chamber of  $\mathcal{P}(\Sigma)$ . Looking at the isomorphism

$$F_{\sigma(\Delta)} = (\mathcal{P}(\sigma(\Delta)) \cap \Delta_0) \xrightarrow{\sim} \mathcal{P}(\Sigma) \cap \Delta = D$$

induced by  $\gamma(\Delta) \in \mathcal{O}^+(L_{10})$ , we see that the set of walls of  $D$  is equal to

$$\{ F_{\Xi}^{\gamma(\Delta)} \mid \Xi \in \mathcal{S} \text{ satisfies } \Sigma \subset \Xi \text{ and } |\Sigma| + 1 = |\Xi| \}.$$

**Definition 3.15.** Let  $w := F_{\Xi}^{\gamma(\Delta)}$  be a wall of the induced chamber  $D$  of  $\mathcal{P}(\Sigma)$ . An induced chamber *adjacent to  $D$  across the wall  $w$*  is the unique induced chamber  $D'$  of  $\mathcal{P}(\Sigma)$  such that  $D \neq D'$  and that  $w$  is a wall of  $D'$ . We say that an induced chamber  $D'$  is *adjacent to  $D$*  if  $D'$  is adjacent to  $D$  across some wall of  $D$ .

**Algorithm 3.16.** Let  $D$  and  $w$  be as in Definition 3.15. By the following method, we can obtain a Vinberg chamber  $\Delta' \in \mathcal{V}_\Sigma$  and an isometry  $\gamma(\Delta') \in \mathcal{O}^+(L_{10})$  such that  $\mathcal{P}(\Sigma) \cap \Delta'$  is the induced chamber adjacent to  $D$  across  $w$ . Let  $\Xi$  be the element of  $\mathcal{S}$  such that  $w = F_{\Xi}^{\gamma(\Delta)}$ , and let  $W(\Xi, L_{10})$  denote the subgroup of  $\mathcal{O}^+(L_{10})$  generated by the reflections  $s_r$  associated with the roots  $r \in \Xi$ . Let  $\xi$  be the longest element of the Coxeter group  $W(\Xi, L_{10}) \cong W(\langle \Xi \rangle)$ , which can be calculated by Algorithm 2.3. Then, if  $\mathcal{U}$  is a sufficiently small neighborhood in  $\mathcal{P}_{10}$  of a general point of the face  $F_{\Xi}$  of  $\Delta_0$ , we have

$$\Delta_0^\xi \cap \mathcal{U} = \{ x \in \mathcal{U} \mid \langle x, r \rangle \leq 0 \text{ for all } r \in \Xi \},$$

that is, the Vinberg chamber  $\Delta_0^\xi$  is opposite to  $\Delta_0$  with respect to the face  $F_{\Xi}$ . Then  $\Delta' := \Delta_0^{\xi\gamma(\Delta)}$  is the desired Vinberg chamber, and we have  $\gamma(\Delta') = \xi\gamma(\Delta)$ .  $\blacksquare$

We present the main algorithm of this section, which is based on the generalized Borcherds method ([5], [6], [22]).

**Algorithm 3.17.** Let an element  $\Sigma$  of  $\mathcal{S}$  be given. This algorithm calculates the set  $\{\Sigma' \in \mathcal{S} \mid \Sigma' \sim \Sigma\}$  and a generating set of the stabilizer subgroup  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  of  $\langle \Sigma \rangle$  in  $\mathcal{O}^+(L_{10})$ . We set

$$\mathcal{D} := [\Delta_0], \quad \gamma(\mathcal{D}) := [\text{id}], \quad \sigma(\mathcal{D}) := [\Sigma], \quad \mathcal{G} := \{\}, \quad i := 0,$$

where  $\text{id}$  is the identity of  $\mathcal{O}^+(L_{10})$ . During the calculation, we have the following:

- (i)  $\mathcal{D}$  is a list  $[\Delta_0, \dots, \Delta_j]$  of elements of  $\mathcal{V}_\Sigma$  such that  $\sigma(\Delta_\mu) \neq \sigma(\Delta_\nu)$  if  $\mu \neq \nu$ , where  $\sigma: \mathcal{V}_\Sigma \rightarrow \mathcal{S}$  is defined by (3.3),
- (ii)  $\gamma(\mathcal{D})$  is the list  $[\gamma(\Delta_0), \dots, \gamma(\Delta_j)]$  of elements of  $\mathcal{O}^+(L_{10})$ ,
- (iii)  $\sigma(\mathcal{D})$  is the list  $[\sigma(\Delta_0), \dots, \sigma(\Delta_j)]$  of distinct elements of  $\mathcal{S}$ , and
- (iv)  $\mathcal{G}$  is a set of elements of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$ .

While  $i + 1 \leq j + 1 = |\mathcal{D}|$ , we execute the following calculation.

- (1) Let  $\Delta_i$  be the  $(i+1)$ st element of  $\mathcal{D}$ . By Algorithm 3.16, we calculate Vinberg chambers  $\Delta^{(1)}, \dots, \Delta^{(k)}$  in  $\mathcal{V}_\Sigma$  such that  $\{\mathcal{P}(\Sigma) \cap \Delta^{(\kappa)} \mid \kappa = 1, \dots, k\}$  is the set of induced chambers in  $\mathcal{P}(\Sigma)$  adjacent to the induced chamber  $\mathcal{P}(\Sigma) \cap \Delta_i$ . Note that, in Algorithm 3.16, we also calculate  $\gamma(\Delta^{(\kappa)}) \in \mathrm{O}^+(L_{10})$  for  $\kappa = 1, \dots, k$ .
- (2) For  $\kappa = 1, \dots, k$ , we calculate  $\Sigma^{(\kappa)} := \sigma(\Delta^{(\kappa)})$  by  $\gamma(\Delta^{(\kappa)})$ .
  - (2-1) If  $\Sigma^{(\kappa)} \notin \sigma(\mathcal{D})$ , then we add  $\Delta^{(\kappa)}$  to  $\mathcal{D}$ ,  $\gamma(\Delta^{(\kappa)})$  to  $\gamma(\mathcal{D})$ , and  $\Sigma^{(\kappa)}$  to  $\sigma(\mathcal{D})$  at the end of each list.
  - (2-2) If  $\Sigma^{(\kappa)}$  appears at the  $(m+1)$ st position of  $\sigma(\mathcal{D})$ , then we have  $\Sigma^{(\kappa)} = \sigma(\Delta_m)$ , where  $\Delta_m \in \mathcal{D}$ . We add  $g := \gamma(\Delta^{(\kappa)})^{-1} \cdot \gamma(\Delta_m)$  to  $\mathcal{G}$ . Note that, since  $\mathcal{P}(\Sigma^{(\kappa)})^{\gamma(\Delta^{(\kappa)})} = \mathcal{P}(\Sigma) = \mathcal{P}(\sigma(\Delta_m))^{\gamma(\Delta_m)}$ , we have  $g \in \mathrm{Stab}(\langle \Sigma \rangle, L_{10})$ .
- (3) We increment  $i$  to  $i+1$ .

Since  $|\mathcal{D}| = |\sigma(\mathcal{D})|$  cannot exceed  $|\mathcal{S}| = 1021$ , this algorithm terminates. When it terminates, we output  $\mathcal{D}$  as  $\mathcal{D}_\Sigma$ ,  $\sigma(\mathcal{D})$  as  $\sigma(\mathcal{D}_\Sigma)$ , and  $\mathcal{G}$  as  $\mathcal{G}_\Sigma$ .  $\blacksquare$

**Proposition 3.18.** *We have  $\sigma(\mathcal{D}_\Sigma) = \{\Sigma' \in \mathcal{S} \mid \Sigma' \sim \Sigma\}$ . Moreover, the finite set  $\mathcal{G}_\Sigma$  together with  $W(\Sigma, L_{10})$  generates  $\mathrm{Stab}(\langle \Sigma \rangle, L_{10})$ .*

*Proof.* Let  $G$  be the subgroup of  $\mathrm{O}^+(L_{10})$  generated by the union of  $\mathcal{G}_\Sigma$  and  $W(\Sigma, L_{10})$ . Note that  $G \subset \mathrm{Stab}(\langle \Sigma \rangle, L_{10})$ , and hence  $\mathcal{P}(\Sigma)^g = \mathcal{P}(\Sigma)$  holds for all  $g \in G$ . First we show that, for any  $\Delta \in \mathcal{V}_\Sigma$ , there exists an element  $g \in G$  such that  $\Delta^g \in \mathcal{D}_\Sigma$ . An *adjacent sequence* is a sequence  $D_{(0)}, \dots, D_{(N)}$  of induced chambers of  $\mathcal{P}(\Sigma)$  such that  $D_{(\nu-1)}$  and  $D_{(\nu)}$  are adjacent for each  $\nu = 1, \dots, N$ . The number  $N$  is called the *length* of the adjacent sequence  $D_{(0)}, \dots, D_{(N)}$ . For  $\Delta \in \mathcal{V}_\Sigma$ , let  $d(\Delta)$  be the minimum of the lengths of adjacent sequences from  $\mathcal{P}(\Sigma) \cap \Delta_0$  to  $\mathcal{P}(\Sigma) \cap \Delta$ . Since  $\mathcal{P}(\Sigma)$  is connected and covered by induced chambers, we have  $d(\Delta) < \infty$  for any  $\Delta \in \mathcal{V}_\Sigma$ . Suppose that

$$\mathcal{B} := \{ \Delta \in \mathcal{V}_\Sigma \mid \Delta^g \notin \mathcal{D}_\Sigma \text{ for any } g \in G \}$$

is non-empty. Let  $\Delta_{\min} \in \mathcal{B}$  be an element such that  $N := d(\Delta_{\min}) \leq d(\Delta)$  holds for any  $\Delta \in \mathcal{B}$ . We put  $\Delta_{(0)} := \Delta_0$  and  $\Delta_{(N)} := \Delta_{\min}$ . Then we have an adjacent sequence

$$\mathcal{P}(\Sigma) \cap \Delta_{(0)}, \mathcal{P}(\Sigma) \cap \Delta_{(1)}, \dots, \mathcal{P}(\Sigma) \cap \Delta_{(N-1)}, \mathcal{P}(\Sigma) \cap \Delta_{(N)}$$

of length  $N$  from  $\mathcal{P}(\Sigma) \cap \Delta_0$  to  $\mathcal{P}(\Sigma) \cap \Delta_{\min}$ . The minimality of  $N = d(\Delta_{\min})$  implies that there exists an element  $g$  of  $G$  such that  $\Delta_{(N-1)}^g \in \mathcal{D}_\Sigma$ . We put  $\Delta_i := \Delta_{(N-1)}^g$ . Since  $\mathcal{P}(\Sigma)^g = \mathcal{P}(\Sigma)$ , we see that  $\mathcal{P}(\Sigma) \cap \Delta_{(N)}^g$  is an induced chamber adjacent to  $\mathcal{P}(\Sigma) \cap \Delta_i$ . Therefore, when Algorithm 3.17 processed  $\Delta_i \in \mathcal{D}$ , there must be a Vinberg chamber  $\Delta^{(\kappa)}$  among  $\Delta^{(1)}, \dots, \Delta^{(k)}$  such that  $\mathcal{P}(\Sigma) \cap \Delta^{(\kappa)} = \mathcal{P}(\Sigma) \cap \Delta_{(N)}^g$ . By Lemma 3.12, we have an element  $g' \in W(\Sigma, L_{10}) \subset G$  such that  $\Delta_{(N)}^{gg'} = \Delta^{(\kappa)}$ . Since  $\Delta_{\min} \in \mathcal{B}$  implies that  $\Delta^{(\kappa)} = \Delta_{\min}^{gg'}$  is not in  $\mathcal{D}_\Sigma$ , the case (2-2) must have occurred. Therefore  $\sigma(\Delta^{(\kappa)}) = \sigma(\Delta_{\min}^{gg'})$  should have been added to  $\sigma(\mathcal{D}_\Sigma)$ , and there exists an element  $\Delta_m \in \mathcal{D}_\Sigma$  such that

$$g'' := \gamma(\Delta^{(\kappa)})^{-1} \cdot \gamma(\Delta_m) \in \mathcal{G}_\Sigma.$$

Then  $gg'g'' \in G$  and  $\Delta_{\min}^{gg'g''} = \Delta_m \in \mathcal{D}_\Sigma$ , which is a contradiction. Thus  $\mathcal{B} = \emptyset$  is proved.

Next we prove  $\sigma(\mathcal{D}_\Sigma) = \{\Sigma' \in \mathcal{S} \mid \Sigma' \sim \Sigma\}$ . Lemma 3.13 implies that all  $\Sigma' \in \sigma(\mathcal{D}_\Sigma)$  satisfies  $\Sigma' \sim \Sigma$ . Suppose that  $\Sigma' \in \mathcal{S}$  satisfies  $\Sigma' \sim \Sigma$ , and let  $h \in \mathrm{O}^+(L_{10})$

be an isometry such that  $\langle \Sigma \rangle = \langle \Sigma' \rangle^h$ . We put  $\Delta := \Delta_0^h$ . Since  $(\mathcal{P}(\Sigma') \cap \Delta_0)^h = \mathcal{P}(\Sigma) \cap \Delta$ , we have  $\Delta \in \mathcal{V}_\Sigma$ . Since  $\mathcal{B} = \emptyset$ , we have an element  $g \in G$  such that  $\Delta^g \in \mathcal{D}_\Sigma$ . Since  $\mathcal{P}(\Sigma)^g = \mathcal{P}(\Sigma)$ , we have  $(\mathcal{P}(\Sigma') \cap \Delta_0)^{hg} = \mathcal{P}(\Sigma) \cap \Delta^g$ , which implies  $\Sigma' = \sigma(\Delta^g)$ . Therefore  $\Sigma'$  belongs to  $\sigma(\mathcal{D}_\Sigma)$ .

We prove  $G \supset \text{Stab}(\langle \Sigma \rangle, L_{10})$ . Let  $h$  be an element of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$ . Then we have  $\Delta_0^h \in \mathcal{V}_\Sigma$ . Since  $\mathcal{B} = \emptyset$ , we have an element  $g \in G$  such that  $\Delta_0^{hg} \in \mathcal{D}_\Sigma$ . Since  $\mathcal{P}(\Sigma)^{hg} = \mathcal{P}(\Sigma)$ , we have  $(\mathcal{P}(\Sigma) \cap \Delta_0)^{hg} = \mathcal{P}(\Sigma) \cap \Delta_0^{hg}$ , and therefore  $\sigma(\Delta_0^{hg}) = \Sigma$ . Since elements of  $\sigma(\mathcal{D}_\Sigma)$  are all distinct, we have  $\Delta_0^{hg} = \Delta_0$  and hence  $h = g^{-1} \in G$  holds.  $\square$

We execute Algorithm 3.17 for all  $\Sigma \in \mathcal{S}$ , and confirm that  $\Sigma \sim \Sigma'$  holds if and only if their ADE-types  $\tau(\Sigma)$  and  $\tau(\Sigma')$  coincide. Hence we obtain the following:

**Theorem 3.19.** *The set  $\mathcal{N}/\mathcal{O}^+(L_{10})$  is identified with the set  $\tau(\mathcal{S})$  of ADE-types of elements of  $\mathcal{S}$ . In particular, we have  $|\mathcal{N}/\mathcal{O}^+(L_{10})| = 86$ .  $\square$*

Thus we have classified, up to the action of  $\mathcal{O}^+(L_{10})$ , all primitive sublattices of  $L_{10}$  that appear as  $\overline{R}_f$  in Theorem 1.3. In order to complete the proof of Theorem 1.3, we now enumerate all ADE-configurations  $\Phi_f$  in each  $\overline{R}_f$ .

**3.2. Stabilizer subgroups.** Note that  $W(\Sigma, L_{10})$  is a normal subgroup of the stabilizer subgroup  $\text{Stab}(\langle \Sigma \rangle, L_{10})$ . We put

$$\text{Stab}(\Sigma, L_{10}) := \{ g \in \mathcal{O}^+(L_{10}) \mid \Sigma^g = \Sigma \},$$

which is a subgroup of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  and is mapped isomorphically to the quotient group  $\text{Stab}(\langle \Sigma \rangle, L_{10})/W(\Sigma, L_{10})$ . Hence the rows of the following commutative diagram are *splitting* exact sequences:

$$(3.5) \quad \begin{array}{ccccccc} 1 & \rightarrow & W(\Sigma, L_{10}) & \rightarrow & \text{Stab}(\langle \Sigma \rangle, L_{10}) & \xrightarrow{\tilde{\kappa}} & \text{Stab}(\Sigma, L_{10}) \rightarrow 1 \\ & & \downarrow \wr & & \downarrow \text{res} & & \downarrow \text{res} \\ 1 & \rightarrow & W(\langle \Sigma \rangle) & \rightarrow & \mathcal{O}(\langle \Sigma \rangle) & \xrightarrow{\kappa} & \text{Aut}(\Sigma) \rightarrow 1, \end{array}$$

where the vertical arrows are the restriction homomorphisms.

**Corollary 3.20.** *The group  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  is generated by the union of  $W(\Sigma, L_{10})$  and  $\tilde{\kappa}(\mathcal{G}_\Sigma)$ , and the group  $\text{Stab}(\Sigma, L_{10})$  is generated by  $\tilde{\kappa}(\mathcal{G}_\Sigma)$ .  $\square$*

We can calculate  $\tilde{\kappa}(\mathcal{G}_\Sigma)$  by the same method as Algorithm 2.2. We can also calculate a generating set  $\kappa(\text{res}(\mathcal{G}_\Sigma)) = \text{res}(\tilde{\kappa}(\mathcal{G}_\Sigma))$  of the subgroup

$$(3.6) \quad H_\Sigma := \text{Im} \left( \text{Stab}(\langle \Sigma \rangle, L_{10}) \xrightarrow{\text{res}} \mathcal{O}(\langle \Sigma \rangle) \xrightarrow{\kappa} \text{Aut}(\Sigma) \right)$$

of  $\text{Aut}(\Sigma)$ .

**3.3. Embeddings of an ADE-configuration.** Let  $\Phi$  be an ADE-configuration with  $|\Phi| < 10$ . Let  $\text{Emb}(\Phi)$  denote the set of all embeddings of  $\Phi$  into  $L_{10}$ . Then  $\text{Aut}(\Phi)$  acts on  $\text{Emb}(\Phi)$  from the left, and  $\mathcal{O}^+(L_{10})$  acts on  $\text{Emb}(\Phi)$  from the right. In this section, we calculate the set  $\text{Aut}(\Phi) \backslash \text{Emb}(\Phi) / \mathcal{O}^+(L_{10})$ , and prove Theorem 1.3.

Let  $f: \Phi \hookrightarrow L_{10}$  be an embedding. We denote by  $\Phi_f$  the image of  $f$ , by  $R_f$  the sublattice of  $L_{10}$  generated by  $\Phi_f$ , and by  $\overline{R}_f$  the primitive closure of  $R_f$  in  $L_{10}$ . Then  $\overline{R}_f$  corresponds to an even overlattice  $R_{(f)}$  of  $\langle \Phi \rangle$  via the isometry  $\langle \Phi \rangle \xrightarrow{\sim} R_{(f)}$  given by  $f$ , and we have  $\overline{R}_f \in \mathcal{N}$  by Lemma 3.8. Recall that  $\tau(\mathcal{S})$  is the set of

ADE-types of all  $\Sigma \in \mathcal{S}$ . We consider the following condition on an even overlattice  $\overline{R}$  of  $\langle \Phi \rangle$ .

(#)  $\overline{R}$  is a root lattice whose ADE-type belongs to  $\tau(\mathcal{S})$ .

Then Theorem 3.19 implies the following:

- (a) For any  $f \in \text{Emb}(\Phi)$ , the even overlattice  $R_{(f)}$  of  $\langle \Phi \rangle$  satisfies (#), and there exist an element  $g \in O^+(L_{10})$  and an element  $\Sigma \in \mathcal{S}$  such that  $\overline{R}_{fg} = \langle \Sigma \rangle$ .
- (b) Suppose that an even overlattice  $\overline{R}$  of  $\langle \Phi \rangle$  satisfies (#). Then there exist an element  $\Sigma \in \mathcal{S}$  and an isometry  $\bar{f}: \overline{R} \xrightarrow{\sim} \langle \Sigma \rangle$ . The restriction of  $\bar{f}$  to  $\Phi \subset \overline{R}$  gives an embedding  $f \in \text{Emb}(\Phi)$ .

Therefore we can calculate  $\text{Aut}(\Phi) \backslash \text{Emb}(\Phi) / O^+(L_{10})$  by the following method.

(1) Let  $\mathcal{L}(\Phi)$  denote the set of even overlattices of  $\langle \Phi \rangle$ , on which  $\text{Aut}(\Phi)$  acts from the right. We calculate the set  $\mathcal{L}(\Phi) / \text{Aut}(\Phi)$  of orbits of this action, and for each orbit  $o \in \mathcal{L}(\Phi) / \text{Aut}(\Phi)$ , we choose a representative  $\overline{R} \in o$  and calculate the stabilizer subgroup  $\text{Stab}(\overline{R}, \Phi)$  of  $\overline{R}$  in the finite group  $\text{Aut}(\Phi)$ .

(2) For  $\overline{R} \in \mathcal{L}(\Phi)$ , let  $[\overline{R}] \in \mathcal{L}(\Phi) / \text{Aut}(\Phi)$  denote the orbit containing  $\overline{R}$ . Similarly, for  $\Sigma \in \mathcal{S}$ , let  $[\Sigma] \in \mathcal{N} / O^+(L_{10})$  denote the orbit containing  $\langle \Sigma \rangle \in \mathcal{N}$ . We define a subset  $\mathcal{I}_\Phi$  of  $\mathcal{L}(\Phi) / \text{Aut}(\Phi) \times \mathcal{N} / O^+(L_{10})$  by

$$\mathcal{I}_\Phi := \{ ([\overline{R}], [\Sigma]) \mid \overline{R} \text{ satisfies (\#) and } \tau(\overline{R}) = \tau(\Sigma) \}.$$

For each pair  $([\overline{R}], [\Sigma]) \in \mathcal{I}_\Phi$ , we define a set  $\overline{\text{emb}}([\overline{R}], [\Sigma])$  as follows. Let  $\text{Isom}(\overline{R}, \langle \Sigma \rangle)$  denote the set of all isomorphisms from  $\overline{R}$  to  $\langle \Sigma \rangle$ , on which  $\text{Stab}(\overline{R}, \Phi)$  acts from the left and  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  acts from the right. We put

$$\overline{\text{emb}}([\overline{R}], [\Sigma]) := \text{Stab}(\overline{R}, \Phi) \backslash \text{Isom}(\overline{R}, \langle \Sigma \rangle) / \text{Stab}(\langle \Sigma \rangle, L_{10}).$$

Then the set  $\text{Aut}(\Phi) \backslash \text{Emb}(\Phi) / O^+(L_{10})$  is the disjoint union of  $\overline{\text{emb}}([\overline{R}], [\Sigma])$ , where  $([\overline{R}], [\Sigma])$  runs through the set  $\mathcal{I}_\Phi$ .

3.3.1. *Execution of task (1).* The task (1) above can be easily carried out by Proposition 2.5. We calculate the set of all totally isotropic subgroups of the discriminant form  $q_{\langle \Phi \rangle}$  up to the action of  $\text{Aut}(\Phi)$ . Executing this calculation for all 157 ADE-configurations  $\Phi$  with  $|\Phi| < 10$ , we obtain the following:

**Theorem 3.21.** *Let  $\Phi$  be an ADE-configuration with  $|\Phi| < 10$ . Suppose that  $\overline{R}_1$  and  $\overline{R}_2$  are even overlattices of  $\langle \Phi \rangle$ . If  $\overline{R}_1$  and  $\overline{R}_2$  are root lattices with the same ADE-type, then there exists an automorphism  $g \in \text{Aut}(\Phi)$  such that  $\overline{R}_1^g = \overline{R}_2$ .  $\square$*

**Example 3.22.** Suppose that  $\Phi = \{r_1, \dots, r_n\}$  is of ADE-type  $nA_1$ . Then the discriminant group  $A_{\langle \Phi \rangle}$  of  $\langle \Phi \rangle$  is an  $n$ -dimensional  $\mathbb{F}_2$ -vector space with basis  $r_i^\vee := -r_i/2 \pmod{\langle \Phi \rangle}$  ( $i = 1, \dots, n$ ), and  $q_{\langle \Phi \rangle}$  is given by

$$q_{\langle \Phi \rangle}(a_1 r_1^\vee + \dots + a_n r_n^\vee) = -(a_1^2 + \dots + a_n^2) / 2 \pmod{2\mathbb{Z}}, \quad \text{where } a_1, \dots, a_n \in \mathbb{F}_2.$$

Hence a subspace  $\mathcal{C}$  of  $A_{\langle \Phi \rangle} = \mathbb{F}_2^n$  is totally isotropic with respect to  $q_{\langle \Phi \rangle}$  if and only if  $\mathcal{C}$  is a doubly-even linear code in  $\mathbb{F}_2^n$ . (See Ebeling [8] for the terminologies on codes.) We classify these codes up to the action of  $\text{Aut}(\Phi) \cong \mathfrak{S}_n$  by a brute-force method. In the case  $n = 8$ , there exist exactly 8 doubly-even linear codes in  $\mathbb{F}_2^8$  up to  $\mathfrak{S}_8$ . The corresponding even overlattices of  $\langle \Phi \rangle$  are given in Table 3.1, where  $[i_1 \dots i_k]$  denotes the codeword  $(r_{i_1}^\vee + \dots + r_{i_k}^\vee) \pmod{\langle \Phi \rangle}$  in  $\mathbb{F}_2^8$ .



$\dim \mathcal{C}$	$\mathcal{C}$	$\tau(\overline{R})$	Condition (#)
0	0	$8A_1$	no
1	$\langle [5678] \rangle$	$4A_1 + D_4$	no
1	$\langle [12 \dots 8] \rangle$	not a root lattice	no
2	$\langle [3678], [4578] \rangle$	$2A_1 + D_6$	no
2	$\langle [1678], [2345] \rangle$	$2D_4$	no
3	$\langle [2678], [3578], [4568] \rangle$	$A_1 + E_7$	yes
3	$\langle [1678], [2578], [3478] \rangle$	$D_8$	yes
4	$\langle [1678], [2578], [3568], [4567] \rangle$	$E_8$	yes

TABLE 3.1. Even overlattices of the root lattice of type  $8A_1$ 

3.3.2. *Execution of task (2).* Suppose that  $([\overline{R}], [\Sigma])$  is an element of  $\mathcal{I}_\Phi$ . We describe a method to calculate the set  $\overline{\text{emb}}([\overline{R}], [\Sigma])$ . We can find an ADE-basis of the even overlattice  $\overline{R}$  by the method described in Section 2.4, and hence we can write an element  $\varphi_0 \in \text{Isom}(\overline{R}, \langle \Sigma \rangle)$  explicitly. Using  $\varphi_0$  as a reference point, we identify  $\text{Isom}(\overline{R}, \langle \Sigma \rangle)$  with  $O(\langle \Sigma \rangle)$ . Moreover, since we have a natural injective homomorphism  $\text{Stab}(\overline{R}, \Phi) \hookrightarrow O(\overline{R})$ , we can regard  $\text{Stab}(\overline{R}, \Phi)$  as a subgroup of  $O(\langle \Sigma \rangle)$ .

*Remark 3.23.* By Algorithm 2.2, we can choose  $\varphi_0 \in \text{Isom}(\overline{R}, \langle \Sigma \rangle)$  in such a way that the image by  $\varphi_0$  of the connected component

$$\{ x \in \overline{R} \otimes \mathbb{R} \mid \langle x, r \rangle > 0 \text{ for all } r \in \Phi \}$$

of  $(\overline{R} \otimes \mathbb{R})^\circ = (\langle \Phi \rangle \otimes \mathbb{R})^\circ$  contains the connected component

$$\{ y \in \langle \Sigma \rangle \otimes \mathbb{R} \mid \langle y, r \rangle > 0 \text{ for all } r \in \Sigma \}$$

of  $(\langle \Sigma \rangle \otimes \mathbb{R})^\circ$ . See Section 2.4.

Let  $H_\Phi$  denote the image of  $\text{Stab}(\overline{R}, \Phi) \subset O(\langle \Sigma \rangle)$  by the quotient homomorphism  $O(\langle \Sigma \rangle) \rightarrow \text{Aut}(\Sigma)$  by  $W(\langle \Sigma \rangle)$ , which can be calculated by Algorithm 2.2. Since  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  contains  $W(\Sigma, L_{10})$ , we see that

$$\overline{\text{emb}}([\overline{R}], [\Sigma]) := \text{Stab}(\overline{R}, \Phi) \backslash O(\langle \Sigma \rangle) / \text{Stab}(\langle \Sigma \rangle, L_{10}) = H_\Phi \backslash \text{Aut}(\Sigma) / H_\Sigma,$$

where  $H_\Sigma$  is defined by (3.6). By this computation, we obtain the following:

**Theorem 3.24.** *The set  $\overline{\text{emb}}([\overline{R}], [\Sigma])$  consists of a single element for any ADE-configuration  $\Phi$  with  $|\Phi| < 10$  and any pair  $([\overline{R}], [\Sigma]) \in \mathcal{I}_\Phi$ .  $\square$*

Now we can prove Theorem 1.3.

*Proof of Theorem 1.3.* The assertion (1) follows from Lemma 3.8. The assertion (2) follows from Theorems 3.19, 3.21 and 3.24.  $\square$

**3.4. The stabilizer subgroup of  $\Phi_f$  in  $O^+(L_{10})$ .** Suppose that  $f \in \text{Emb}(\Phi)$  and  $\Sigma \in \mathcal{S}$  satisfy  $\bar{R}_f = \langle \Sigma \rangle$ , and that  $f$  induces an element  $\varphi_0 \in \text{Isom}(\bar{R}, \langle \Sigma \rangle)$  satisfying the condition in Remark 3.23. We put

$$\text{Stab}(\Phi_f, L_{10}) := \{ g \in O^+(L_{10}) \mid \Phi_f^g = \Phi_f \}.$$

It is obvious that  $\text{Stab}(\Phi_f, L_{10}) \subset \text{Stab}(\langle \Sigma \rangle, L_{10})$ . The purpose of this section is to calculate a finite generating set  $\mathcal{G}'_{\Phi}$  of  $\text{Stab}(\Phi_f, L_{10})$ . This set  $\mathcal{G}'_{\Phi}$  will be used in the next section for the classifications of strongly equivalence classes of RDP-Enriques surfaces.

The following general algorithm is used several times in this section.

**Algorithm 3.25.** Let  $G$  be a group generated by  $\gamma_1, \dots, \gamma_N \in G$ . Suppose that  $G$  acts on a finite set  $S$ , and let  $s_0$  be an element of  $S$ . This algorithm calculates a set  $\mathcal{R}_0$  of elements of  $G$  such that  $g \mapsto s_0^g$  gives a bijection from  $\mathcal{R}_0$  to the orbit  $s_0^G := \{s_0^g \mid g \in G\}$  of  $s_0$  under the action of  $G$ . This algorithm also calculates a finite generating set  $\mathcal{G}_0$  of the stabilizer subgroup

$$\text{Stab}(s_0, G) := \{ g \in G \mid s_0^g = s_0 \}.$$

We set  $\Gamma := \{\gamma_1, \dots, \gamma_N, \gamma_1^{-1}, \dots, \gamma_N^{-1}\}$ . We then put  $h_0 := \text{id} \in G$ , and

$$\mathcal{R} := [h_0], \quad \mathcal{O} := [s_0], \quad \mathcal{G} := \{ \}, \quad i := 0.$$

During the calculation, we have the following:

- (i)  $\mathcal{R}$  is a list  $[h_0, h_1, \dots, h_j]$  of elements of  $G$ , and  $\mathcal{O}$  is the list  $[s_0, s_1, \dots, s_j]$  of *distinct* elements of  $s_0^G$  such that  $s_\mu = s_0^{h_\mu}$  holds for  $\mu = 0, \dots, j$ , and
- (ii)  $\mathcal{G}$  is a set of elements of  $\text{Stab}(s_0, G)$ .

While  $i + 1 \leq j + 1 = |\mathcal{R}| = |\mathcal{O}|$ , we execute the following calculation.

- (1) Let  $h_i$  be the  $(i + 1)$ st element of  $\mathcal{R}$ , and let  $s_i = s_0^{h_i}$  be the  $(i + 1)$ st element of  $\mathcal{O}$ . For each  $\gamma \in \Gamma$ , we execute the following:
  - (1-1) If  $s_i^\gamma = s_0^{h_i \gamma} \notin \mathcal{O}$ , then we add  $h_i \gamma$  to  $\mathcal{R}$  and  $s_i^\gamma$  to  $\mathcal{O}$  at the end of each list, whereas
  - (1-2) if  $s_i^\gamma = s_0^{h_i \gamma}$  is equal to the  $(m + 1)$ st element  $s_m = s_0^{h_m}$  of  $\mathcal{O}$ , then we add  $h_i \gamma h_m^{-1} \in \text{Stab}(s_0, G)$  to  $\mathcal{G}$ .
- (2) We increment  $i$  to  $i + 1$ .

Since  $|\mathcal{O}| = |\mathcal{R}|$  cannot exceed  $|S|$ , this algorithm terminates. When it terminates, it outputs  $\mathcal{R}$  as  $\mathcal{R}_0$ ,  $\mathcal{O}$  as  $\mathcal{O}_0$ , and  $\mathcal{G}$  as  $\mathcal{G}_0$ . ■

**Proposition 3.26.** *The set  $\mathcal{O}_0$  is equal to  $s_0^G$ , and  $\mathcal{G}_0$  generates  $\text{Stab}(s_0, G)$ .*

*Proof.* The proof is similar to the proof of Proposition 3.18. The details are left to the reader. □

*Remark 3.27.* We have calculated a finite generating set  $\tilde{\kappa}(\mathcal{G}_\Sigma) \cup W(\Sigma, L_{10})$  of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$ , and hence we can obtain a finite generating set of  $\text{Stab}(\Phi_f, L_{10})$  by applying Algorithm 3.25 to the action of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  on the set of ADE-configurations of roots in  $\langle \Sigma \rangle$ . This method, however, takes too much time and memory for many  $f \in \text{Emb}(\Phi)$ . Therefore we use the following method.

We put

$$V := \langle \Sigma \rangle \otimes \mathbb{R} = \langle \Phi_f \rangle \otimes \mathbb{R}.$$

Then  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  acts on  $V$  via the restriction homomorphism  $\text{res}$  (see (3.5)). We denote the action of  $g \in \text{Stab}(\langle \Sigma \rangle, L_{10})$  on  $V$  simply by  $v \mapsto v^g$  without writing  $\text{res}$ . For an ADE-configuration  $\Psi$  of roots of  $\langle \Sigma \rangle$ , we put

$$\Gamma(\Psi) := \{ y \in V \mid \langle y, r \rangle \geq 0 \text{ for all } r \in \Psi \}.$$

By the assumption on  $f$  (see Remark 3.23), we have

$$\Gamma(\Sigma) \subset \Gamma(\Phi_f).$$

We calculate the finite set

$$\mathcal{W}(\Phi_f, \Sigma) := \{ w \in W(\Sigma, L_{10}) \mid \Gamma(\Sigma)^w \subset \Gamma(\Phi_f) \}.$$

Recall that  $\tilde{\kappa}(\mathcal{G}_\Sigma)$  is a generating set of the subgroup  $\text{Stab}(\Sigma, L_{10})$  of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$ . Therefore, by Algorithm 3.25, we can calculate a finite subset

$$\mathcal{R}_{\Phi_f, \Sigma} = \{h_1, \dots, h_N\}$$

of  $\text{Stab}(\Sigma, L_{10})$  such that  $g \mapsto \Phi_f^g$  gives a bijection from  $\mathcal{R}_{\Phi_f, \Sigma}$  to the orbit

$$\mathcal{O}(\Phi_f) := \{ \Phi_f^g \mid g \in \text{Stab}(\Sigma, L_{10}) \}.$$

We set  $\mathcal{G}' := \{ \}$ . When the calculation below terminates, this set  $\mathcal{G}'$  is the desired generating set  $\mathcal{G}'_\Phi$  of  $\text{Stab}(\Phi_f, L_{10})$ . Recall that the upper row of (3.5) is a splitting exact sequence. Hence each element  $g$  of  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  is uniquely written as  $\tilde{\kappa}(g)w$ , where  $w \in W(\Sigma, L_{10})$ . We consider the coset decomposition

$$\text{Stab}(\langle \Sigma \rangle, L_{10}) = \bigsqcup_{w \in W(\Sigma, L_{10})} \text{Stab}(\Sigma, L_{10}) \cdot w,$$

and for each  $w \in W(\Sigma, L_{10})$ , we consider the set

$$\Xi(w) := (\text{Stab}(\Sigma, L_{10}) \cdot w) \cap \text{Stab}(\Phi_f, L_{10}).$$

If  $w \notin \mathcal{W}(\Phi_f, \Sigma)$ , then we obviously have  $\Xi(w) = \emptyset$ . Therefore we assume that  $w \in \mathcal{W}(\Phi_f, \Sigma)$ . We then calculate the image  $\mathcal{O}(\Phi_f)^w$  of the orbit  $\mathcal{O}(\Phi_f)$  by  $w$ . If  $\mathcal{O}(\Phi_f)^w$  does not contain  $\Phi_f$ , then we have  $\Xi(w) = \emptyset$ . Therefore we assume that there exists an element  $h_i$  of  $\mathcal{R}_{\Phi_f, \Sigma}$  such that  $\Phi_f^{h_i w} = \Phi_f$ . We add  $h_i w$  in  $\mathcal{G}'$ . Every element of  $\Xi(w)$  is written uniquely as  $h_i g w$ , where  $g \in \text{Stab}(\Sigma, L_{10})$ . Since  $h_i g w = h_i w (w^{-1} g w)$ , an element  $g \in \text{Stab}(\Sigma, L_{10})$  satisfies  $h_i g w \in \Xi(w)$  if and only if  $(\Phi_f^{w^{-1}})^g = \Phi_f^{w^{-1}}$ . Using the finite generating set  $\tilde{\kappa}(\mathcal{G}_\Sigma)$  of  $\text{Stab}(\Sigma, L_{10})$  and Algorithm 3.25, we calculate a finite generating set  $\mathcal{G}''(w)$  of

$$\text{Stab}(\Phi_f^{w^{-1}}, \Sigma, L_{10}) := \{ g \in \text{Stab}(\Sigma, L_{10}) \mid (\Phi_f^{w^{-1}})^g = \Phi_f^{w^{-1}} \}.$$

We then append  $w^{-1} \mathcal{G}''(w) w$  to  $\mathcal{G}'$ . Thus a finite generating set  $\mathcal{G}'_\Phi$  of  $\text{Stab}(\Phi_f, L_{10})$  is computed.

The group  $\text{Stab}(\Phi_f, L_{10})$  is, in general, infinite. However we have examples as follows.

**Example 3.28.** We consider the case where  $\tau(\Phi) = 8A_1$  and  $\tau(\Sigma) = E_8$  (no. 88 of Table 1.1). In this case, the method above still takes too much computation time, but we can calculate  $\text{Stab}(\Phi_f, L_{10})$  as follows. Since  $R_f = \langle \Sigma \rangle$  is unimodular,  $L_{10}$  is the orthogonal direct-sum of  $\langle \Sigma \rangle$  and a hyperbolic plane  $U$ . Hence  $\text{Stab}(\langle \Sigma \rangle, L_{10})$  is contained in the subgroup  $\text{O}(\langle \Sigma \rangle) \times \text{O}^+(U)$  of  $\text{O}^+(L_{10})$ . Note that  $\text{O}^+(U)$  is of

order 2. Let  $g_U$  be the non-trivial element of  $O^+(U)$ . Then the kernel of the natural homomorphism

$$\text{Stab}(\Phi_f, L_{10}) \rightarrow \text{Aut}(\Phi)$$

is generated by  $(\text{id}, g_U) \in O(\langle \Sigma \rangle) \times O^+(U)$ , and the image of this homomorphism is  $\{\sigma \in \text{Aut}(\Phi) \cong \mathfrak{S}_8 \mid \mathcal{C}^\sigma = \mathcal{C}\}$ , where  $\mathcal{C}$  is the doubly-even linear code in  $A_{\langle \Phi \rangle} \cong \mathbb{F}_2^8$  corresponding to the even overlattice  $\bar{R} \cong \langle \Sigma \rangle$ , that is,  $\mathcal{C}$  is an extended Hamming code [8, Chapter 1]. Therefore we have  $|\text{Stab}(\Phi_f, L_{10})| = 2688$ , and we can easily obtain a generating set of  $\text{Stab}(\Phi_f, L_{10})$ .

The case where  $\tau(\Phi) = 9A_1$  and  $\tau(\Sigma) = A_1 + E_8$  (no. 146 of Table 1.1) is also treated in the similar method. We have  $|\text{Stab}(\Phi_f, L_{10})| = 1344$  in this case.

#### 4. GEOMETRIC REALIZABILITY

Let  $\Phi$  be an ADE-configuration with  $|\Phi| < 10$ . An embedding  $f: \Phi \hookrightarrow L_{10}$  is said to be *geometrically realized by an RDP-Enriques surface*  $(Y, \rho)$  if there exists an isometry  $S_Y \xrightarrow{\sim} L_{10}$  that maps  $\mathcal{P}_Y$  to  $\mathcal{P}_{10}$  and  $\Phi_\rho$  to  $\Phi_f$  bijectively. If an embedding  $f': \Phi' \hookrightarrow L_{10}$  is equivalent (in the sense of Definition 3.1) to a geometrically realizable embedding  $f: \Phi \hookrightarrow L_{10}$ , then  $f'$  is also geometrically realizable. The purpose of this section is to introduce a lattice  $\bar{M}_f$  corresponding to the lattice  $\bar{M}_\rho$  associated with an RDP-Enriques surface  $(Y, \rho)$ , and give a criterion for the geometric realizability. By means of these tools, we classify the strong equivalence classes of RDP-Enriques surfaces, and prove Theorem 1.7.

For a lattice  $L$ , let  $L(2)$  denote the lattice obtained from  $L$  by multiplying the intersection form by 2. The orthogonal direct-sum of lattices  $L$  and  $L'$  is denoted by  $L \oplus L'$ .

**4.1. Geometry of an Enriques involution.** An involution  $\varepsilon: X \rightarrow X$  of a  $K3$  surface  $X$  is said to be an *Enriques involution* if  $\varepsilon$  has no fixed points, or equivalently, the quotient  $X/\langle \varepsilon \rangle$  is an Enriques surface. The following follows from the classification of 2-elementary  $K3$  surfaces due to Nikulin [15].

**Proposition 4.1** (Nikulin). *Let  $X$  be a  $K3$  surface, and let  $\varepsilon: X \rightarrow X$  be an involution, which acts on  $H^2(X, \mathbb{Z})$  from the right by the pull-back. Suppose that  $\varepsilon$  acts on  $H^0(X, \Omega_X^2)$  as the multiplication by  $-1$ . Let  $S_X^+$  denote the primitive sublattice  $\{v \in S_X \mid v^\varepsilon = v\}$  of  $S_X$ . Then  $\varepsilon$  is an Enriques involution if and only if  $S_X^+$  is isomorphic to  $L_{10}(2)$ .  $\square$*

Let  $\varepsilon: X \rightarrow X$  be an Enriques involution, and let  $\pi: X \rightarrow Y$  denote the universal covering of the Enriques surface  $Y := X/\langle \varepsilon \rangle$ . Then  $S_Y$  is isomorphic to  $L_{10}$ , and  $\pi^*: S_Y \rightarrow S_X$  induces an isometry  $S_Y(2) \cong \pi^* S_Y = S_X^+$ . We also have  $\pi^{*-1}(\mathcal{P}_X) = \mathcal{P}_Y$ . By an isometry  $S_Y \cong L_{10}$  that maps  $\mathcal{P}_Y$  to  $\mathcal{P}_{10}$ , the notion of Vinberg chambers in  $\mathcal{P}_Y$  makes sense. We also use the notation (3.1) and (3.2) for  $S_Y$ . We put

$$\begin{aligned} \text{Nef}(X) &:= \{x \in \mathcal{P}_X \mid \langle x, C' \rangle \geq 0 \text{ for all curves } C' \subset X\}, \\ \text{Nef}(Y) &:= \{x \in \mathcal{P}_Y \mid \langle x, C \rangle \geq 0 \text{ for all curves } C \subset Y\}. \end{aligned}$$

It is well-known that  $\text{Nef}(X)$  is a standard fundamental domain of the action of  $W(S_X)$  on  $\mathcal{P}_X$ . It is obvious that  $\pi^{*-1}(\text{Nef}(X)) = \text{Nef}(Y)$ . Since  $\text{Nef}(Y)$  is bounded by the hyperplanes  $([C])^\perp$  of  $\mathcal{P}_Y$  perpendicular to the classes of smooth rational curves  $C$  on  $Y$ , and a smooth rational curve on  $Y$  has self-intersection number  $-2$ , the cone  $\text{Nef}(Y)$  is tessellated by Vinberg chambers.

In the following, we fix an ample class  $a \in S_Y$  such that  $\langle r, a \rangle \neq 0$  for any root  $r$  of  $S_Y$ . For example, we choose an interior point of a Vinberg chamber contained in  $\text{Nef}(Y)$ . In particular, if  $R$  is a negative-definite root sublattice of  $S_Y$ , then  $\langle -, a \rangle$  is an element of  $\text{Hom}(R, \mathbb{R})^\circ$  in the notation of Section 2.4. Since  $\pi^*a$  is ample on  $X$ , we have  $\langle r', \pi^*a \rangle \neq 0$  for any root  $r'$  of  $S_X$ . We put

$$N_Y := \{ v \in S_X \mid \langle v, y \rangle = 0 \text{ for all } y \in \pi^*S_Y \}.$$

If  $N_Y$  contained a root, then there would be an effective divisor of  $X$  contracted by  $\pi$ , which is absurd. Hence we have the following:

**Lemma 4.2.** *The negative-definite even lattice  $N_Y$  contains no roots.*  $\square$

Note that  $\varepsilon$  acts on  $N_Y$  as the multiplication by  $-1$ . We put

$$\mathcal{T}_Y := \{ t \in N_Y \mid \langle t, t \rangle = -4 \}.$$

If  $r \in \text{Roots}(S_Y)$  and  $t \in \mathcal{T}_Y$ , then  $(\pi^*r + t)/2 \in S_X \otimes \mathbb{Q}$  is of square-norm  $-2$ . A root  $r$  of  $S_Y$  is said to be *liftable* if  $r' := (\pi^*r + t)/2$  is contained in  $S_X$  for some  $t \in \mathcal{T}_Y$ , and when this is the case, the root  $r'$  of  $S_X$  is called a *lift* of  $r$ .

**Lemma 4.3.** *For a liftable root  $r \in S_Y$ , there exist exactly two lifts  $r'$  and  $r''$  of  $r$ , and they satisfy  $\pi^*r = r' + r''$ ,  $r'' = r'^\varepsilon$ , and  $\langle r', r'' \rangle = 0$ .*

*Proof.* If  $r' = (\pi^*r + t)/2$  with  $t \in \mathcal{T}_Y$  is a lift of  $r$ , then so is  $r'' := (\pi^*r - t)/2$ . Suppose that  $t' \in \mathcal{T}_Y$  gives a lift  $(\pi^*r + t')/2$  of  $r$ . Then we have  $(t - t')/2 \in N_Y$ . By Lemma 4.2, we see that  $\langle t, t' \rangle = -4$  or  $\langle t, t' \rangle \geq 4$ . By the Cauchy-Schwarz inequality for  $N_Y$ , we have  $|\langle t, t' \rangle| \leq 4$ . Therefore we have  $t' = \pm t$ .  $\square$

An effective divisor  $D$  of  $Y$  is said to be *splitting* if there exists an effective divisor  $D'$  of  $X$  such that  $\pi^*(D) = D' + \varepsilon(D')$  and  $\langle D', \varepsilon(D') \rangle = 0$ . Note that an effective divisor of  $Y$  is splitting if each connected component of its support is simply connected. In particular, a smooth rational curve on  $Y$  is splitting.

**Lemma 4.4.** *Let  $r$  be a root of  $S_Y$  such that  $\langle r, a \rangle > 0$ . Then  $r$  is the class of a splitting effective divisor of  $Y$  if and only if  $r$  is liftable.*

*Proof.* Suppose that  $r = [D]$ , where  $D$  is a splitting effective divisor, and suppose that  $\pi^*(D) = D' + \varepsilon(D')$  with  $\langle D', \varepsilon(D') \rangle = 0$ . Then  $t := [D'] - [\varepsilon(D')]$  belongs to  $\mathcal{T}_Y$  and  $[D'] = (\pi^*r + t)/2 \in S_X$ . Therefore  $r$  is liftable. Conversely, suppose that  $r$  has lifts  $r'$  and  $r'' = r'^\varepsilon$ . Since  $r'$  is a root of  $S_X$  and  $\langle r', \pi^*a \rangle > 0$ , there exists an effective divisor  $D'$  of  $X$  such that  $r' = [D']$ . Then we have  $\pi^*r = r' + r''$ ,  $r'' = [\varepsilon(D')]$  and  $\langle D', \varepsilon(D') \rangle = 0$  by Lemma 4.3. Let  $D$  be the effective divisor of  $Y$  such that  $\pi^*(D) = D' + \varepsilon(D')$ . Then  $D$  is splitting and we have  $r = [D]$ .  $\square$

Let  $H$  be a nef divisor of  $Y$  such that  $\langle H, H \rangle > 0$ , and let  $h \in \text{Nef}(Y) \cap S_Y$  be the class of  $H$ . If  $k$  is a sufficiently large and divisible integer, then the complete linear system  $|kH|$  is base-point free and the Stein factorization of the morphism  $Y \rightarrow \mathbb{P}^N$  induced by  $|kH|$  gives rise to an RDP-Enriques surface

$$\rho_h : Y \rightarrow \bar{Y}.$$

We calculate the set  $\Phi_h := \Phi_{\rho_h}$  of the classes of smooth rational curves contracted by  $\rho_h$ . Let  $[h]^\perp$  denote the orthogonal complement in  $S_Y$  of the sublattice  $[h] := \mathbb{Z}h$  generated by  $h$ . Then  $[h]^\perp$  is negative-definite. We put

$$(4.1) \quad \Pi_h^+ := \{ r \in \text{Roots}([h]^\perp) \mid \langle r, a \rangle > 0 \}, \quad L_h^+ := \{ r \in \Pi_h^+ \mid r \text{ is liftable} \},$$

and consider the root sublattice  $\langle L_h^+ \rangle$  of  $[h]^\perp$  generated by  $L_h^+$  (possibly of rank 0).

**Proposition 4.5.** *The ADE-configuration  $\Phi_h$  of the RDP-Enriques surface  $(Y, \rho_h)$  is equal to the ADE-basis of  $\langle L_h^+ \rangle$  associated with the linear form  $\langle -, a \rangle$  by the correspondence described in Section 2.4.*

*Proof.* By Lemma 4.4, every element  $r$  of  $L_h^+$  is the class of a splitting effective divisor  $D_r$  contracted by  $\rho_h$ . Since the class of any smooth rational curve contracted by  $\rho_h$  is in  $L_h^+$ , the divisor  $D_r$  is irreducible if and only if  $r \in L_h^+$  is indecomposable in  $L_h^+$  in the sense of Definition 2.4. Therefore  $\Phi_h$  is equal to the set  $\Phi'_h$  of indecomposable elements of  $L_h^+$ . In particular, the set  $\Phi'_h$  is an ADE-configuration of roots of  $S_Y$ , and the vectors of  $\Phi'_h$  are linearly independent. Since every element of  $L_h^+$  is a linear combination of the indecomposable elements, we have  $\langle \Phi'_h \rangle = \langle L_h^+ \rangle$ . Since the ADE-basis of the root lattice  $\langle L_h^+ \rangle$  associated with the linear form  $\langle -, a \rangle \in \text{Hom}(\langle L_h^+ \rangle, \mathbb{R})^\circ$  is unique, we obtain the proof.  $\square$

**4.2. Lattices associated with an RDP-Enriques surface.** Let  $(Y, \rho)$  be an RDP-Enriques surface, and  $a \in S_Y$  an ample class such that  $\langle a, r \rangle \neq 0$  holds for any  $r \in \text{Roots}(S_Y)$ . Let  $C_1, \dots, C_n$  be the smooth rational curves on  $Y$  contracted by  $\rho: Y \rightarrow \bar{Y}$ , so that  $\Phi_\rho = \{[C_1], \dots, [C_n]\}$ . Since the Dynkin diagram of  $\Phi_\rho$  is a disjoint union of trees, we have the following:

**Lemma 4.6.** *Let  $D$  be an effective divisor of  $Y$  contracted by  $\rho$ . Then each connected component of the support of  $D$  is simply connected. In particular,  $D$  is splitting.*  $\square$

Let  $\pi: X \rightarrow Y$  be the universal covering, and let  $C'_i$  and  $C''_i$  be the two connected components of  $\pi^{-1}(C_i)$ . Lemma 4.6 implies that, interchanging  $C'_i$  and  $C''_i$  if necessary, we can assume that

$$(4.2) \quad \langle C_i, C_j \rangle = \langle C'_i, C'_j \rangle = \langle C''_i, C''_j \rangle, \quad \text{and} \quad \langle C'_i, C''_j \rangle = 0$$

hold for all  $i, j$ . We put

$$\phi([C_i]) := [C'_i] - [C''_i] \in N_Y.$$

Note that  $[C'_i]$  and  $[C''_i]$  are the lifts of  $[C_i]$  associated with  $\phi([C_i]) \in \mathcal{T}_Y$ .

*Remark 4.7.* Let  $c$  be the number of connected components of the Dynkin diagram of  $\Phi_\rho$ . Then there exist exactly  $2^c$  possibilities for the choice of the map  $\phi$ .

By (4.2), we have

$$(4.3) \quad \langle \phi([C_i]), \phi([C_j]) \rangle = 2\langle [C_i], [C_j] \rangle,$$

and hence  $\phi$  defines an embedding

$$(4.4) \quad \phi: \langle \Phi_\rho \rangle(2) \hookrightarrow N_Y.$$

Recall that  $R_\rho$  is the sublattice of  $S_Y$  generated by  $\Phi_\rho$ , and that  $\bar{R}_\rho$  is the primitive closure of  $R_\rho$  in  $S_Y$ . Recall also that  $\Phi_\rho^\circ$  is the subset  $\{[C'_1], [C''_1], \dots, [C'_n], [C''_n]\}$  of  $\text{Roots}(S_X)$ , that  $M_\rho$  is the sublattice of  $S_X$  generated by  $\pi^*S_Y$  and  $\Phi_\rho^\circ$ , and that  $\bar{M}_\rho$  is the primitive closure of  $M_\rho$  in  $S_X$ . Then  $M_\rho$  is an even overlattice of the orthogonal direct-sum

$$B_\rho := \pi^*S_Y \oplus \text{Im } \phi.$$

Thus we obtain a sequence of inclusions

$$(4.5) \quad \pi^*\Phi_\rho \subset \pi^*R_\rho \subset \pi^*\bar{R}_\rho \subset \pi^*S_Y \subset B_\rho \subset M_\rho \subset \bar{M}_\rho.$$

**Lemma 4.8.** *We have  $\Phi_\rho^\sim = \{ r' \in \text{Roots}(\overline{M}_\rho) \mid r' + r'^\varepsilon \in \pi^* \Phi_\rho \}$ .*

*Proof.* The inclusion  $\subset$  is obvious. Suppose that  $r' \in \text{Roots}(\overline{M}_\rho)$  satisfies  $r' + r'^\varepsilon = \pi^* r$  for some  $r \in \Phi_\rho$ . Since  $\langle r, a \rangle > 0$ , we have  $\langle r', \pi^* a \rangle > 0$ , and hence we have an effective divisor  $D'$  of  $X$  such that  $r' = [D']$ . Let  $D$  be the effective divisor of  $Y$  such that  $\pi^* D = D' + \varepsilon(D')$ . Then  $r = [D] \in \Phi_\rho$  and hence  $D = C_i$  for some  $i$ . Therefore  $r' \in \Phi_\rho^\sim$ .  $\square$

We can now prove Lemma 1.6 stated in the introduction.

*Proof of Lemma 1.6.* Since  $\pi^* S_Y$  (resp.  $\pi'^* S_{Y'}$ ) is the invariant sublattice of the Enriques involution on  $X$  (resp. on  $X'$ ), the strong equivalence isometry  $\mu$  is compatible with the action of the Enriques involutions on  $\overline{M}_\rho$  and  $\overline{M}_{\rho'}$ . Since  $\mu_Y$  maps  $\Phi_\rho$  to  $\Phi_{\rho'}$  bijectively, Lemma 4.8 implies that  $\mu$  maps  $\Phi_\rho^\sim$  to  $\Phi_{\rho'}^\sim$  bijectively.  $\square$

**Proposition 4.9.** *The set  $\{r \in \text{Roots}(\overline{R}_\rho) \mid r \text{ is liftable}\}$  is equal to  $\text{Roots}(R_\rho)$ .*

*Proof.* Let  $h_\rho \in S_Y$  be the class of the pull-back by  $\rho: Y \rightarrow \overline{Y}$  of an ample divisor of  $\overline{Y}$ . Since  $\Phi_\rho$  is orthogonal to  $h_\rho$ , the sublattices  $R_\rho$  and  $\overline{R}_\rho$  of  $S_Y$  are orthogonal to  $h_\rho$ . Suppose that  $r \in \text{Roots}(\overline{R}_\rho)$  is liftable. Replacing  $r$  by  $-r$  if necessary, we can assume that  $\langle r, a \rangle > 0$ . By Lemma 4.4, the root  $r$  is the class of a splitting effective divisor  $D$ . Since  $\langle r, h_\rho \rangle = 0$ , the divisor  $D$  is contracted by  $\rho$ . Therefore  $D$  is a linear combination of  $C_1, \dots, C_n$ . Thus we have  $r \in \text{Roots}(R_\rho)$ . Suppose that  $r \in \text{Roots}(R_\rho)$ . Replacing  $r$  by  $-r$  if necessary, we can assume that  $\langle r, a \rangle > 0$ , and hence there exists an effective divisor  $D = \sum a_i C_i$  such that  $r = [D]$ . Then  $D$  is splitting by Lemma 4.6, and hence  $r$  is liftable by Lemma 4.4.  $\square$

**4.3. A criterion for geometric realizability.** For an embedding  $f: \Phi \hookrightarrow L_{10}$  of an ADE-configuration  $\Phi = \{r_1, \dots, r_n\}$  into  $L_{10}$ , we will construct a sequence in (4.7), which is a lattice-theoretic counterpart of (4.5). For  $r_i \in \Phi$ , we denote by  $r_i^+ \in \Phi_f$  the image of  $r_i$  by  $f$ , so that  $\Phi_f = \{r_1^+, \dots, r_n^+\}$ . Let  $\Phi^- = \{r_1^-, \dots, r_n^-\}$  be a copy of  $\Phi$  with a fixed isomorphism  $\Phi \cong \Phi^-$  denoted by  $r_i \mapsto r_i^-$ . We put

$$B_\Phi := L_{10}(2) \oplus \langle \Phi^- \rangle(2).$$

We denote by

$$\varpi^*: L_{10} \xrightarrow{\sim} L_{10}(2), \quad \varphi: \langle \Phi^- \rangle \xrightarrow{\sim} \langle \Phi^- \rangle(2),$$

the identity maps on the underlying  $\mathbb{Z}$ -modules. Hence we have

$$(4.6) \quad \langle \varpi^* r_i^+, \varpi^* r_j^+ \rangle = \langle \varphi(r_i^-), \varphi(r_j^-) \rangle = 2\langle r_i, r_j \rangle.$$

We define vectors  $r'_i$  and  $r''_i$  of  $B_\Phi^\vee \subset B_\Phi \otimes \mathbb{Q}$  by

$$r'_i := (\varpi^* r_i^+ + \varphi(r_i^-))/2, \quad r''_i := (\varpi^* r_i^+ - \varphi(r_i^-))/2.$$

Let  $M_f$  denote the submodule of  $B_\Phi^\vee$  generated by  $B_\Phi$  and  $r'_1, \dots, r'_n$ . Then  $M_f$  is an even overlattice of  $B_\Phi$ . Note that  $r''_i$  also belongs to  $M_f$ . Let  $\overline{M}_f$  be an even overlattice of  $M_f$ . Recall that  $R_f$  is the sublattice of  $L_{10}$  generated by  $\Phi_f$ , and that  $\overline{R}_f$  is the primitive closure of  $R_f$  in  $L_{10}$ . Then we have a sequence of inclusions

$$(4.7) \quad \varpi^* \Phi_f \subset \varpi^* R_f \subset \varpi^* \overline{R}_f \subset \varpi^* L_{10} \subset B_\Phi \subset M_f \subset \overline{M}_f.$$

Let  $N(\overline{M}_f)$  denote the orthogonal complement of  $\varpi^* L_{10}$  in  $\overline{M}_f$ . We put

$$\mathcal{T}(\overline{M}_f) := \{ t \in N(\overline{M}_f) \mid \langle t, t \rangle = -4 \}.$$

A root  $r$  of  $\overline{R}_f$  is said to be  $\overline{M}_f$ -*liftable* if there exists an element  $t \in \mathcal{T}(\overline{M}_f)$  such that  $r' := (\varpi^*r + t)/2 \in \overline{M}_f \otimes \mathbb{Q}$  is contained in  $\overline{M}_f$ , and when this is the case, we say that  $r'$  is an  $\overline{M}_f$ -*lift* of  $r$ .

**Lemma 4.10.** *Every root of  $R_f$  is  $\overline{M}_f$ -liftable for any even overlattice  $\overline{M}_f$  of  $M_f$ .*

*Proof.* Let  $r^+$  be a root of  $R_f$ . We can write  $r^+$  as  $\sum a_i r_i^+$  with  $a_i \in \mathbb{Z}$ . Then, putting  $r^- := \sum a_i r_i^- \in \langle \Phi^- \rangle$ , we have  $\varphi(r^-) \in \mathcal{T}(M_f) \subset \mathcal{T}(\overline{M}_f)$ , and hence  $r^+$  has an  $\overline{M}_f$ -lift  $(\varpi^*r^+ + \varphi(r^-))/2 = \sum a_i r_i'$ .  $\square$

**Definition 4.11.** We define the following conditions (C1)-(C4) on  $\overline{M}_f$ . Let  $L_{K3}$  denote the  $K3$  lattice, that is,  $L_{K3}$  is an even unimodular lattice of signature  $(3, 19)$ , which is unique up to isomorphism.

- (C1) The lattice  $\overline{M}_f$  can be embedded primitively into  $L_{K3}$ .
- (C2) The sublattice  $\varpi^*L_{10}$  is primitive in  $\overline{M}_f$ .
- (C3) The negative-definite even lattice  $N(\overline{M}_f)$  contains no roots.
- (C4) The set  $\{r \in \text{Roots}(\overline{R}_f) \mid r \text{ is } \overline{M}_f\text{-liftable}\}$  is equal to  $\text{Roots}(R_f)$ .

**Definition 4.12.** We say that an even overlattice  $\overline{M}_f$  of  $M_f$  is *strongly realized* by an RDP-Enriques surface  $(Y, \rho)$  if there exists an isometry

$$m: \overline{M}_f \xrightarrow{\sim} \overline{M}_\rho$$

with the following properties; the isometry  $m$  maps  $\varpi^*L_{10}$  to  $\pi^*S_Y$  isomorphically, and the isometry  $m_Y: L_{10} \xrightarrow{\sim} S_Y$  induced by  $m$  maps  $\mathcal{P}_{10}$  to  $\mathcal{P}_Y$  and  $\Phi_f$  to  $\Phi_\rho$  bijectively. An isometry  $m: \overline{M}_f \xrightarrow{\sim} \overline{M}_\rho$  satisfying these conditions is called a *strong-realization isometry*.

*Remark 4.13.* If an even overlattice  $\overline{M}_f$  of  $M_f$  is strongly realized by an RDP-Enriques surface  $(Y, \rho)$ , then  $f: \Phi \hookrightarrow L_{10}$  is geometrically realized by  $(Y, \rho)$ .

**Theorem 4.14.** *If an embedding  $f: \Phi \hookrightarrow L_{10}$  is geometrically realized by an RDP-Enriques surface  $(Y, \rho)$ , then there exist an even overlattice  $\overline{M}_f$  of  $M_f$  strongly realized by  $(Y, \rho)$ .*

*Proof.* By the assumption, we have an isometry  $m_Y: L_{10} \xrightarrow{\sim} S_Y$  that maps  $\mathcal{P}_{10}$  to  $\mathcal{P}_Y$  and  $\Phi_f$  to  $\Phi_\rho$  bijectively. We use the notation about  $(Y, \rho)$  fixed in Section 4.2, and index the elements  $r_1, \dots, r_n$  of  $\Phi$  in such a way that  $m_Y(r_i^+) = [C_i]$  holds for  $i = 1, \dots, n$ . Then  $m_Y$  induces an isometry

$$m^+ : \varpi^*L_{10} \xrightarrow{\sim} \pi^*S_Y$$

that maps  $\varpi^*r_i^+$  to  $[\pi^*(C_i)]$ . By (4.3), we also have an isometry

$$m^- : \langle \Phi^- \rangle(2) \xrightarrow{\sim} \text{Im}(\phi: \langle \Phi_\rho \rangle(2) \hookrightarrow N_Y)$$

that maps  $\varphi(r_i^-)$  to  $\phi([C_i]) = [C_i'] - [C_i'']$ . Thus we have an isometry

$$m := m^+ \oplus m^- : B_\Phi \xrightarrow{\sim} B_\rho.$$

Then  $m \otimes \mathbb{Q}$  maps  $r_i'$  to  $[C_i']$  and  $r_i''$  to  $[C_i'']$ . Therefore we obtain an isometry  $m: M_f \xrightarrow{\sim} M_\rho$ . Let  $\overline{M}_f$  be the even overlattice of  $M_f$  corresponding to the even overlattice  $\overline{M}_\rho$  of  $M_\rho$  via  $m$ , so that  $m: M_f \xrightarrow{\sim} M_\rho$  extends to  $m: \overline{M}_f \xrightarrow{\sim} \overline{M}_\rho$ . Then  $m$  induces an isomorphism from the sequence (4.7) to the sequence (4.5). In particular,  $m$  is a strong-realization isometry.  $\square$



**Theorem 4.15.** *If an even overlattice  $\overline{M}_f$  of  $M_f$  is strongly realized by an RDP-Enriques surface  $(Y, \rho)$ , then  $\overline{M}_f$  satisfies (C1)-(C4).*

*Proof.* Let  $m: \overline{M}_f \xrightarrow{\sim} \overline{M}_\rho$  be a strong-realization isometry. Since  $\overline{M}_\rho$  is a primitive sublattice of the primitive sublattice  $S_X$  of the  $K3$ -lattice  $H^2(X, \mathbb{Z}) \cong L_{K3}$ , the lattice  $\overline{M}_f \cong \overline{M}_\rho$  satisfies (C1). Since  $\pi^*S_Y$  is primitive in  $S_X$  and hence in  $\overline{M}_\rho$ , the lattice  $\overline{M}_f$  satisfies (C2). Since the isometry  $m$  maps the lattice  $N(\overline{M}_f)$  isomorphically to a sublattice of  $N_Y$ , Lemma 4.2 implies that  $\overline{M}_f$  satisfies (C3). If  $r \in \text{Roots}(\overline{R}_f)$  is  $\overline{M}_f$ -liftable, then  $m_Y(r) \in \text{Roots}(\overline{R}_\rho)$  is liftable, because  $m$  maps  $\mathcal{T}(\overline{M}_f)$  to a subset of  $\mathcal{T}_Y$ . Hence the isometry  $m_Y: L_{10} \xrightarrow{\sim} S_Y$  induced by  $m$  induces the horizontal injection in the commutative diagram below:

$$\begin{array}{ccc} \{r \in \text{Roots}(\overline{R}_f) \mid r \text{ is } \overline{M}_f\text{-liftable}\} & \hookrightarrow & \{r \in \text{Roots}(\overline{R}_\rho) \mid r \text{ is liftable}\} \\ \text{by Lemma 4.10 } \uparrow & & \Downarrow \text{ by Proposition 4.9} \\ \text{Roots}(R_f) & \xrightarrow[\text{by } m_Y]{\sim} & \text{Roots}(R_\rho). \end{array}$$

By Proposition 4.9 and Lemma 4.10, we see that the upward injection in the left column of the diagram above is a bijection. Hence  $\overline{M}_f$  satisfies (C4).  $\square$

**Theorem 4.16.** *Let  $\overline{M}_f$  be an even overlattice of  $M_f$  satisfying (C1)-(C4). Then there exists an RDP-Enriques surface  $(Y, \rho)$  that strongly realizes  $\overline{M}_f$ .*

*Proof.* The main tool is the Torelli theorem for  $K3$  surfaces and the surjectivity of the period map of  $K3$  surfaces (see [4, Chapter VIII]). By (C1) for  $\overline{M}_f$  and the surjectivity of the period map, there exists a  $K3$  surface  $X$  equipped with a marking

$$\mu: \overline{M}_f \cong S_X.$$

Our task is to construct an Enriques involution  $\varepsilon$  on  $X$ , and an RDP-Enriques surface  $\rho: Y := X/\langle \varepsilon \rangle \rightarrow \overline{Y}$  that strongly realizes  $\overline{M}_f$ .

Replacing  $\mu$  by  $-\mu$  if necessary, we can assume that  $\mu$  induces an embedding

$$\mu \circ \varpi^*|_{\mathcal{P}} : \mathcal{P}_{10} \hookrightarrow \mathcal{P}_X.$$

We put  $\mathcal{P}(\Phi_f) := (\Phi_f)^\perp = ([\Phi_f]^\perp \otimes \mathbb{R}) \cap \mathcal{P}_{10}$ , which is a positive cone of the hyperbolic sublattice  $[\Phi_f]^\perp = R_f^\perp = \overline{R}_f^\perp$  of  $L_{10}$ . There exists a Vinberg chamber  $\Delta$  of  $L_{10}$  such that  $\mathcal{P}(\Phi_f) \cap \Delta$  contains a non-empty open subset of  $\mathcal{P}(\Phi_f)$ . Moving  $\Delta$  by an element of the subgroup  $W(\Phi_f, L_{10}) \subset O^+(L_{10})$  generated by the reflections  $s_r$  with respect to the roots  $r \in \Phi_f$ , we can assume that  $\Delta$  satisfies the following:

$$(4.8) \quad \langle r, v \rangle \geq 0 \quad \text{holds for all } r \in \Phi_f \text{ and } v \in \Delta.$$

Let  $r'$  be a root of  $\overline{M}_f$ . By (C3) for  $\overline{M}_f$ , the locus

$$(r')^\perp := \{x \in \mathcal{P}_{10} \mid \langle \varpi^*x, r' \rangle = 0\}$$

is a hyperplane of  $\mathcal{P}_{10}$ . Note that the family  $\{(r')^\perp \mid r' \in \text{Roots}(\overline{M}_f)\}$  of hyperplanes of  $\mathcal{P}_{10}$  is locally finite. Let  $\eta'$  be a general point of  $\mathcal{P}(\Phi_f) \cap \Delta \cap (L_{10} \otimes \mathbb{Q})$ , let  $\mathcal{U}$  be a sufficiently small neighborhood of  $\eta'$  in  $\mathcal{P}_{10}$ , and let  $\alpha'$  be a general point of  $\mathcal{U} \cap \Delta \cap (L_{10} \otimes \mathbb{Q})$ . We have a positive integer  $c$  such that  $\alpha := c\alpha' \in L_{10}$  and  $\eta := c\eta' \in L_{10}$ . Then we have the following:

- (i) there exist no roots  $r'$  in  $\overline{M}_f$  such that  $\varpi^*\alpha \in (r')^\perp$ ,
- (ii) there exist no roots  $r'$  in  $\overline{M}_f$  such that  $\langle \varpi^*\alpha, r' \rangle > 0$  and  $\langle \varpi^*\eta, r' \rangle < 0$ , and
- (iii) the set of  $r \in \text{Roots}(L_{10})$  satisfying  $\langle \eta, r \rangle = 0$  is equal to  $\text{Roots}(\overline{R}_f)$ .

We then put

$$a_X := \mu \circ \varpi^*|_{\mathcal{P}}(\alpha) \in \mathcal{P}_X \cap S_X, \quad h_X := \mu \circ \varpi^*|_{\mathcal{P}}(\eta) \in \mathcal{P}_X \cap S_X.$$

By (i), we compose the marking  $\mu$  by an element of  $W(S_X)$  and assume that  $a_X$  is ample. Then (ii) implies that  $h_X$  is nef.

By (C2), the sublattice  $\mu(\varpi^*L_{10})$  of  $S_X$  is primitive in  $S_X$  and hence is primitive in the even unimodular lattice  $H^2(X, \mathbb{Z})$ . Let  $K$  denote the orthogonal complement of  $\mu(\varpi^*L_{10})$  in  $H^2(X, \mathbb{Z})$ . Then we have an isomorphism

$$A_K \cong A_{\mu(\varpi^*L_{10})} \cong A_{L_{10}(2)} \cong \mathbb{F}_2^{10}$$

of discriminant groups by [14]. Since  $A_K$  is 2-elementary, the scalar multiplication by  $-1$  on  $K$  acts on  $A_K$  trivially. By [14] again, we obtain an isometry  $\varepsilon'$  of  $H^2(X, \mathbb{Z})$  that preserves each of  $\mu(\varpi^*L_{10})$  and  $K$ , induces the identity on  $\mu(\varpi^*L_{10})$ , and induces the scalar multiplication by  $-1$  on  $K$ . Since  $\varepsilon'$  acts on  $H^0(X, \Omega_X^2)$  as the scalar multiplication by  $-1$ , it preserves the Hodge structure of  $H^2(X, \mathbb{Z})$ . In particular, the action of  $\varepsilon'$  preserves  $S_X$ . Since  $\varepsilon'$  fixes the ample class  $a_X$ , the action of  $\varepsilon'$  preserves the nef cone of  $X$ . Hence the Torelli theorem implies that  $\varepsilon'$  comes from an involution  $\varepsilon: X \rightarrow X$ . By Proposition 4.1, we see that  $\varepsilon$  is an Enriques involution. Let  $\pi: X \rightarrow Y := X/\langle \varepsilon \rangle$  be the quotient morphism. Since  $\mu(\varpi^*L_{10})$  is the invariant sublattice of  $\varepsilon'$  in  $H^2(X, \mathbb{Z})$ , we have  $\mu(\varpi^*L_{10}) = \pi^*S_Y$ . Therefore  $\mu$  induces an isometry

$$\mu_Y: L_{10} \xrightarrow{\sim} S_Y.$$

Note that  $a := \mu_Y(\alpha)$  is ample on  $Y$  and  $h := \mu_Y(\eta)$  is nef on  $Y$ , because  $\pi^*a = a_X$  and  $\pi^*(h) = h_X$ . Let  $H$  be the nef divisor of  $Y$  whose class is  $h$ . We construct a birational morphism  $\rho_h: Y \rightarrow \bar{Y}$  by  $H$  as in Section 4.1. It remains to show that  $\mu_Y$  maps  $\Phi_f$  to  $\Phi_h := \Phi_{\rho_h}$  bijectively. Recall the definition (4.1) of  $\Pi_h^+$ . The property (iii) of  $\eta$  above implies that  $\mu_Y$  maps  $\text{Roots}(\bar{R}_f)$  to  $\Pi_h^+ \cup (-1)\Pi_h^+$  bijectively. Hence  $\mu_Y$  identifies  $\bar{R}_f$  and the sublattice  $\langle \Pi_h^+ \rangle$  of  $S_Y$  generated by  $\Pi_h^+$ . Since  $\mu$  is an isometry from  $\bar{M}_f$  to  $S_X$ , the map  $\mu$  induces a bijection from  $\mathcal{T}(\bar{M}_f)$  to  $\mathcal{T}_Y$ . Therefore the isometry  $\mu_Y$  induces a bijection from the set of  $\bar{M}_f$ -liftable roots of  $\bar{R}_f$  to the set of liftable root of  $\langle \Pi_h^+ \rangle$ . Recall that  $\alpha$  is in the interior of the Vinberg chamber  $\Delta$ . By (C4) for  $\bar{M}_f$  and (4.8), the ADE-basis of the root sublattice generated by  $\{r \in \text{Roots}(\bar{R}_f) \mid r \text{ is } \bar{M}_f\text{-liftable}\}$  associated with the linear form  $\langle -, \alpha \rangle$  is  $\Phi_f$ . On the other hand, Proposition 4.5 implies that the ADE-basis of the root sublattice generated by  $\{r \in \text{Roots}(\langle \Pi_h^+ \rangle) \mid r \text{ is liftable}\}$  associated with  $\langle -, a \rangle$  is  $\Phi_h$ . Therefore  $\mu_Y$  maps  $\Phi_f$  to  $\Phi_h$  bijectively.  $\square$

**Corollary 4.17.** *An embedding  $f: \Phi \hookrightarrow L_{10}$  is geometrically realizable if and only if there exists an even overlattice  $\bar{M}_f$  of  $M_f$  satisfying the conditions (C1)-(C4).  $\square$*

**Corollary 4.18.** *There exists a bijection between the set of strong equivalence classes of RDP-Enriques surfaces geometrically realizing  $f: \Phi \hookrightarrow L_{10}$  and the set of orbits of the action of the group*

$$U(M_f) := \{ g \in \text{O}^+(M_f) \mid \varpi^*\Phi_f^g = \varpi^*\Phi_f, \varpi^*L_{10}^g = \varpi^*L_{10} \}.$$

*on the set of even overlattices of  $M_f$  satisfying (C1)-(C4).  $\square$*

*Remark 4.19.* Let  $\overline{M}_f$  and  $\overline{M}'_f$  be even overlattices of  $M_f$  such that  $\overline{M}_f \subset \overline{M}'_f$ . If  $\overline{M}'_f$  satisfies (C2), (C3), and (C4), then so does  $\overline{M}_f$ . In particular, if an even overlattice  $\overline{M}_f$  satisfies (C2), (C3), and (C4), then so does  $M_f$ .

A finite generating set of the group  $U(M_f)$  is calculated as follows. First we define a homomorphism

$$\text{Stab}(\Phi_f, L_{10}) \rightarrow \text{O}^+(M_f), \quad g \mapsto \tilde{g}.$$

Let  $g$  be an element of  $\text{Stab}(\Phi_f, L_{10})$ . Then  $g$  induces an automorphism of the ADE-configuration  $\Phi_f$ . Since  $\Phi_f$  and  $\Phi^-$  are canonically isomorphic by  $r_i^+ \mapsto r_i^-$ , we obtain an automorphism  $g^- \in \text{Aut}(\Phi^-)$  and hence an isometry  $g^- \in \text{O}(\langle \Phi^- \rangle)$ . The action of  $g \oplus g^- \in \text{O}(B_\Phi)$  induces a permutation of  $r'_1, \dots, r'_n \in B_\Phi^\vee$ , and hence preserves the even overlattice  $M_f$  of  $B_\Phi$ . Therefore we obtain  $\tilde{g} \in \text{O}^+(M_f)$ .

Let  $c$  be the number of connected components of the Dynkin diagram of  $\Phi^-$ , and let

$$\langle \Phi^- \rangle(2) = \langle \Phi_1^- \rangle(2) \oplus \dots \oplus \langle \Phi_c^- \rangle(2)$$

be the orthogonal direct-sum decomposition of  $\langle \Phi^- \rangle(2)$  according to the connected components of the Dynkin diagram of  $\Phi^-$ . For  $k = 1, \dots, c$ , let  $u'_k \in \text{O}(\langle \Phi^- \rangle(2))$  denote the isometry that acts on the direct-summand  $\langle \Phi_j^- \rangle(2)$  as the identity for  $j \neq k$  and as the multiplication by  $-1$  for  $j = k$ . We then define  $u_k \in \text{O}(B_\Phi)$  to be the direct sum of  $\text{id}_{L_{10}(2)}$  and  $u'_k$ . Then  $u_k$  acts on each of the subsets  $\{r'_j, r''_j\}$  of  $B_\Phi^\vee$ , and hence we can regard  $u_k$  as an element of  $\text{O}^+(M_f)$ .

**Proposition 4.20.** *Suppose that there exists an RDP-Enriques surfaces geometrically realizing  $f: \Phi \hookrightarrow L_{10}$ . Then the group  $U(M_f)$  is generated by the image of the homomorphism  $\text{Stab}(\Phi_f, L_{10}) \rightarrow \text{O}^+(M_f)$  above and the isometries  $u_1, \dots, u_c$ .*

*Proof.* Let  $U' \subset \text{O}^+(M_f)$  be the group generated by the image of the homomorphism  $\text{Stab}(\Phi_f, L_{10}) \rightarrow \text{O}^+(M_f)$  and the isometries  $u_1, \dots, u_c$ . By construction, we have  $U' \subset U(M_f)$ . We prove  $U' \supset U(M_f)$ . Since  $\varpi^* L_{10} = L_{10}(2)$ , we have a natural restriction map

$$\xi: U(M_f) \rightarrow \text{O}^+(\varpi^* L_{10}) = \text{O}^+(L_{10}), \quad g \mapsto g|_{\varpi^* L_{10}}.$$

The image of  $\xi$  is contained in  $\text{Stab}(\Phi_f, L_{10})$  by definition, and  $\xi$  has a section over  $\text{Stab}(\Phi_f, L_{10}) \subset \text{O}^+(L_{10})$ . Therefore it suffices to show that an arbitrary element  $g$  of  $\text{Ker } \xi$  belongs to the subgroup generated by  $u_1, \dots, u_c$ . By the assumption and Remark 4.19, the orthogonal complement  $N(M_f)$  of  $\varpi^* L_{10}$  in  $M_f$  contains no roots. Hence the same argument as in the proof of Lemma 4.3 implies that each  $r_i^+ \in \Phi_f$  has exactly two  $N(M_f)$ -lifts  $r'_i, r''_i$ . Since  $r_i^{+g} = r_i^+$ , we have  $\{r'_i, r''_i\}^g = \{r'_i, r''_i\}$ . Since  $\varphi(r_i^-) = r'_i - r''_i$ , we have  $\varphi(r_i^-)^g = \pm \varphi(r_i^-)$ . By (4.6), if  $\langle r_i, r_j \rangle \neq 0$ , then  $\varphi(r_i^-)^g$  determines  $\varphi(r_j^-)^g$ . Therefore the action of  $g$  on  $\langle \Phi^- \rangle(2)$  is equal to the action of an element of  $\langle u_1, \dots, u_c \rangle$  on  $\langle \Phi^- \rangle(2)$ .  $\square$

**4.4. Complete list of RDP-Enriques surfaces.** In Section 3, we have calculated the complete list of equivalence classes of embeddings  $f: \Phi \hookrightarrow L_{10}$ . From each equivalence class, we choose a representative  $f: \Phi \hookrightarrow L_{10}$ , calculate a finite generating set of the group  $\text{Stab}(\Phi_f, L_{10})$  by the method given in Section 3.4, and calculate the lattice  $M_f$  and a finite generating set of the group  $U(M_f) \subset \text{O}^+(M_f)$ . Then we calculate the finite set  $\mathcal{L}'(M_f)$  of even overlattices of  $M_f$  that satisfy (C2), (C3), (C4) by Proposition 2.5.

*Remark 4.21.* We enumerate even overlattices of  $M_f$  by enlarging successively the corresponding totally isotropic subgroups of the discriminant form  $q_{M_f}$  of  $M_f$ . By Remark 4.19, if  $\overline{M}_f$  fails to satisfy (C2), (C3), or (C4), then we do not have to enlarge the totally isotropic subgroup  $\overline{M}_f/M_f$  any more.

We then decompose  $\mathcal{L}'(M_f)$  into the union of the orbits under the action of  $U(M_f)$ . Note that the image of  $U(M_f)$  by the natural homomorphism  $O(M_f) \rightarrow O(q_{M_f})$  is of course finite, and is calculated from the finite generating set of  $U(M_f)$ . For each orbit  $o$ , we choose a representative element  $\overline{M}_f \in o$ , and check whether  $\overline{M}_f$  satisfies (C1) or not by Proposition 2.6 and Remark 2.7. If  $\overline{M}_f$  satisfies (C1), then  $o$  corresponds to a strong equivalence class. Thus we obtain the complete list of strong equivalence classes of RDP-Enriques surfaces. The result is given in Table 1.1, and Theorem 1.7 is proved.

#### REFERENCES

- [1] Ayşegül Akyol and Alex Degtyarev. Geography of irreducible plane sextics. *Proc. Lond. Math. Soc.* (3), 111(6):1307–1337, 2015.
- [2] Enrique Artal-Bartolo. Sur les couples de Zariski. *J. Algebraic Geom.*, 3(2):223–247, 1994.
- [3] W. Barth and C. Peters. Automorphisms of Enriques surfaces. *Invent. Math.*, 73(3):383–411, 1983.
- [4] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, second edition, 2004.
- [5] Richard Borcherds. Automorphism groups of Lorentzian lattices. *J. Algebra*, 111(1):133–153, 1987.
- [6] Richard E. Borcherds. Coxeter groups, Lorentzian lattices, and  $K3$  surfaces. *Internat. Math. Res. Notices*, 1998(19):1011–1031, 1998.
- [7] J. H. Conway and N. J. A. Sloane. *Sphere packings, lattices and groups*, volume 290 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, New York, third edition, 1999.
- [8] Wolfgang Ebeling. *Lattices and codes*. Advanced Lectures in Mathematics. Springer Spektrum, Wiesbaden, third edition, 2013.
- [9] The GAP Group. GAP - Groups, Algorithms, and Programming. Version 4.7.9; 2015 (<http://www.gap-system.org>).
- [10] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [11] Dong Seon Hwang, Jong Hae Keum, and Hisanori Ohashi. Gorenstein  $\mathbb{Q}$ -homology projective planes. *Sci. China Math.*, 58(3):501–512, 2015.
- [12] J. Keum and D.-Q. Zhang. Fundamental groups of open  $K3$  surfaces, Enriques surfaces and Fano 3-folds. *J. Pure Appl. Algebra*, 170(1):67–91, 2002.
- [13] Rick Miranda and David R. Morrison. *Embeddings of integral quadratic forms*. electronic, 2009, <http://www.math.ucsb.edu/drm/manuscripts/eiqf.pdf>.
- [14] V. V. Nikulin. Integer symmetric bilinear forms and some of their geometric applications. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43(1):111–177, 238, 1979. English translation: *Math USSR-Izv.* 14 (1979), no. 1, 103–167 (1980).
- [15] V. V. Nikulin. Quotient-groups of groups of automorphisms of hyperbolic forms of subgroups generated by 2-reflections. *Dokl. Akad. Nauk SSSR*, 248(6):1307–1309, 1979. English translation: “On factor groups of the automorphism groups of hyperbolic forms modulo subgroups generated by 2-reflections”, *Soviet Math. Dokl.* 20 (1979), no. 5, 1156–1158 (1980).
- [16] V. V. Nikulin. Description of automorphism groups of Enriques surfaces. *Dokl. Akad. Nauk SSSR*, 277(6):1324–1327, 1984. English translation: *Soviet Math. Dokl.* 30 (1984), no. 1, 282–285.
- [17] Sławomir Rams and Matthias Schütt. On Enriques surfaces with four cusps. *Publ. Res. Inst. Math. Sci.*, 54(3):433–468, 2018.

- [18] Matthias Schütt. Moduli of Gorenstein  $\mathbb{Q}$ -homology projective planes. preprint, 2015. arXiv:1505.04163.
- [19] Ichiro Shimada. On normal  $K3$  surfaces. *Michigan Math. J.*, 55(2):395–416, 2007.
- [20] Ichiro Shimada. Lattice Zariski  $k$ -ples of plane sextic curves and  $Z$ -splitting curves for double plane sextics. *Michigan Math. J.*, 59(3):621–665, 2010.
- [21] Ichiro Shimada. Topology of curves on a surface and lattice-theoretic invariants of coverings of the surface. In *Algebraic geometry in East Asia—Seoul 2008*, volume 60 of *Adv. Stud. Pure Math.*, pages 361–382. Math. Soc. Japan, Tokyo, 2010.
- [22] Ichiro Shimada. An algorithm to compute automorphism groups of  $K3$  surfaces and an application to singular  $K3$  surfaces. *Int. Math. Res. Not. IMRN*, (22):11961–12014, 2015.
- [23] Ichiro Shimada. Connected components of the moduli of elliptic  $K3$  surfaces. *Michigan Math. J.*, 67(3):511–559, 2018.
- [24] Ichiro Shimada. On an Enriques surface associated with a quartic Hessian surface. *Canad. J. Math.*, 71(1):213–246, 2019.
- [25] Ichiro Shimada. Rational double points on Enriques surfaces: computational data, 2017. <http://www.math.sci.hiroshima-u.ac.jp/~shimada/K3.html>.
- [26] È. B. Vinberg. Some arithmetical discrete groups in Lobačevskii spaces. In *Discrete subgroups of Lie groups and applications to moduli (Internat. Colloq., Bombay, 1973)*, pages 323–348. Oxford Univ. Press, Bombay, 1975.
- [27] Jin-Gen Yang. Sextic curves with simple singularities. *Tohoku Math. J. (2)*, 48(2):203–227, 1996.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, HIROSHIMA UNIVERSITY, 1-3-1 KAGAMIYAMA, HIGASHI-HIROSHIMA, 739-8526 JAPAN  
*Email address:* `ichiro-shimada@hiroshima-u.ac.jp`